

Treatment of Plaque Psoriasis Involving High-Impact Sites With Icotrokinra, a Targeted Oral Peptide: Pooled Analyses of 4 Phase 3 Placebo-Controlled Trials

Jennifer Soung,¹ April W. Armstrong,² Ronald B. Vender,³ Josep Riera-Monroig,⁴ Michael Sebastian,⁵ Ofelia Reyes-Servin,⁶ Bassey Effiom Edem,⁷ Ming-Chun Hsu,⁶ Shu Li,⁶ Ya-Wen Yang,⁶ Linda Stein Gold⁸

¹Southern California Clinical Research, Santa Ana, and Harbor University of California Los Angeles, CA, USA; ²Department of Dermatology, University of California Los Angeles, Los Angeles, CA, USA; ³McMaster University and Dermatrials Research Inc., Hamilton, ON, Canada; ⁴Dermatology Department, Hospital Clinic de Barcelona, University of Barcelona, Spain; ⁵Dermatological Practice Dr. med. Michael Sebastian, Mahlow, Germany; ⁶Johnson & Johnson, Spring House/Horsham, PA, USA; ⁷Johnson & Johnson, Leiden, The Netherlands; ⁸Henry Ford Health System, West Bloomfield, MI, USA

This presentation was sponsored by Johnson & Johnson.

Presented at: AAD Annual Meeting; March 27-31, 2026; Denver, Colorado, USA.

Disclosures

JS: Served as a speaker, consultant, advisory board member and/or investigator for AbbVie, Amgen, Arcutis, Aslan, Bristol Myers Squibb, Coval Biopharma, Dermavant, Eli Lilly, Johnson & Johnson, KoBio Labs, National Psoriasis Foundation, Novartis, Ortho Dermatologic, Oruka, Pfizer, Regeneron/Sanofi, and UCB.

AWA: Served as a speaker, consultant, and/or investigator for AbbVie, Amgen, Arcutis, Bristol Myers Squibb, Dermavant Sciences, Eli Lilly and Company, Galderma, Incyte, Johnson & Johnson, LEO Pharma, Novartis, Pfizer, Regeneron, Sanofi, Takeda, and UCB.

RBV: Received grants/research support, speakers bureau/honoraria from: AbbVie, Alumis, Amgen, Arcutis, Bausch, Boehringer Ingelheim, Bristol Myers Squibb, Celltrion, Dermavant, Dermira, DICE, Galderma, Incyte, JAMP, Johnson & Johnson, LEO, Lilly, Meiji, Nimbus, Novartis, Organon, Oruka, Pfizer, Sandoz, Sanofi, Sun, Takeda, UCB, and Zai.

JR-M: Participated in clinical trials/received payments for presentations and consultancy/grants to assist with congresses from AbbVie, Almirall, Amgen, Boehringer Ingelheim, Bristol Myers Squibb, Chiesi, Johnson & Johnson, Krystal Biotech, LEO Pharma, Lilly, Novartis, and UCB.

MS: Collaborated with the following companies: AbbVie, Affibody, Allergan, Almirall, Amgen, AstraZeneca, August Wolff, Boehringer Ingelheim, Bristol Myers Squibb, Dermapharm, Dermira, Incyte, Ipsen, Johnson & Johnson, LEO Pharma, Lilly, MedImmune, Menlo Therapeutics, Moonlake, MSD, Mundipharma, Novartis, Pfizer, Regeneron, Sanofi Genzyme, Takeda, UCB Pharma.

OR-S, BEE, M-CH, SL, and Y-WY: Employees of Johnson & Johnson; may own stock/stock options in Johnson & Johnson.

LSG: Served as an investigator, advisor and/or speaker for AbbVie, Amgen, Bristol Myers Squibb, Galderma, Johnson & Johnson, LEO Pharma, Lilly, Pfizer, Regeneron, Sanofi, Takeda.

Background



Plaque psoriasis (PsO) on high-impact sites (e.g., scalp, genitals, hands/feet, nails) can be difficult-to-treat and negatively impact patients' quality of life¹


Patients with high-impact site PsO are candidates for systemic therapy, regardless of their body surface area (BSA) affected, per the International Psoriasis Council consensus statement²



Icetrokinra (ICO) is the first and only targeted oral peptide:

- Precisely blocks the interleukin (IL)-23 receptor and inhibits IL-23 pathway signaling³
- Evaluated for the treatment of moderate-to-severe plaque PsO in the phase 3 ICONIC-LEAD, ICONIC-ADVANCE 1, and ICONIC-ADVANCE 2 studies, and for plaque PsO involving high-impact sites in the phase 3 ICONIC-TOTAL study; primary/co-primary endpoints of each study were met, with no safety signals identified⁴⁻⁶
- In adults and adolescents with high-impact site PsO involvement, ICO demonstrated:
 - Significantly higher scalp and genital PsO clearance rates vs placebo (PBO) at Week (W)16 (ICONIC-TOTAL)⁶
 - Higher rates of scalp, genital, and hand/foot PsO clearance and substantially improved nail PsO vs PBO at W16 (ICONIC-LEAD)⁷

Icetrokinra Blocks IL-23 From Binding to its Receptor



IFN=interferon, IL-12Rβ1=interleukin-12 receptor beta 1, IL-23Ri=interleukin-23 receptor inhibitor.

Objective



Evaluate ICO effects in pooled cohorts of participants (pts) with PsO involving high-impact sites, including the scalp, genitals, hands/feet, and/or nails, across 4 phase 3 studies

This material is distributed for scientific purposes on J&J Medical Connect and is not for promotional use

Pooled ICONIC-LEAD, -TOTAL, -ADVANCE 1 & 2: ICO vs PBO Through W16

Moderate-to-Severe or High-Impact Site Plaque PsO (N=1866*)

Key inclusion criteria

- ≥12 yrs ICONIC-LEAD & -TOTAL
- ≥18 yrs ICONIC-ADVANCE 1 & 2
- Plaque PsO ≥26 weeks
- ICONIC-LEAD, -ADVANCE 1 & 2:
 - BSA ≥10%; PASI score ≥12; IGA score ≥3
 - Candidate for phototherapy or systemic treatment for plaque PsO
- ICONIC-TOTAL:
 - BSA ≥1% and IGA score ≥2
 - At least moderate high-impact PsO involving ≥1 site: ss-IGA score ≥3; sPGA-G score ≥3; hf-PGA score ≥3
 - Candidate for phototherapy or systemic treatment for plaque PsO and failed ≥1 topical

Pts randomized to ICO or PBO from W0-W16 across 4 pooled phase 3 studies:


ICONIC-LEAD⁴
(N=684^b)
2:1

ICONIC-TOTAL⁶
(N=311^c)
2:1

ICONIC-ADVANCE 1⁵
(N=467)
2:1

ICONIC-ADVANCE 2⁵
(N=404)
4:1

(R)



Current Analysis

*Pooled Full Analysis Set^a
(ICO N=1297; PBO N=569)

^aIncludes pooled ICO- and PBO-randomized pts from the phase 3 ICONIC-LEAD, ICONIC-TOTAL, ICONIC-ADVANCE 1, and ICONIC-ADVANCE 2 studies. ^bIncludes 66 adolescents. ^cIncludes 6 adolescents. **hf-PGA**=Physician's Global Assessment of hands and feet, **IGA**=Investigator's Global Assessment, **PASI**=Psoriasis Area and Severity Index, **QD**=once daily, **R**=randomization, **sPGA-G**=static Physician's Global Assessment of Genitalia, **ss-IGA**=scalp-specific IGA.

Outcomes & Analyses

Response Rates Through W24

- Scalp PsO: ss-IGA score 0/1 & 0^{a,c,e}
- Genital PsO: sPGA-G score 0/1 & 0^{a,c,e}
- Hand/foot PsO: hf-PGA score 0/1 & 0^{a,c,e}

Percent Improvement From Baseline Through W24

- Nail PsO: Modified Nail Psoriasis Severity Index (mNAPSI)^{b,d,f}

AEs Through W16


- Safety assessments for pooled ICO and PBO groups

^aAmong pts with a score of ≥ 3 (at least moderate) on ss-IGA, sPGA-G, and hf-PGA, respectively, at baseline. ^bAmong pts with a score of >0 on mNAPSI at baseline.

^cNonresponder imputation / ^dNo improvement from baseline assigned after pts discontinued study drug due to a lack of efficacy or an adverse event (AE) of worsening PsO, or initiated prohibited medication that could impact PsO. Observed data were used for pts who discontinued study drug for other reasons. ^eAfter accounting for the intercurrent events, pts with missing data were considered nonresponders. ^fThe remaining missing data were not imputed.

The prevalence and severity of high-impact site PsO at baseline were balanced between groups

High-Impact Site PsO Among All Pooled Randomized Pts^a (ICO: N=1297 / PBO: N=569)



Severity of High-Impact Site PsO^b

	ICO	PBO
ss-IGA score ≥3, N	935	408
Moderate (3)	80%	77%
Severe (4)	20%	23%
sPGA-G score ≥3, N	334	139
Moderate (3)	75%	68%
Severe (4)	23%	29%
Very severe (5)	2%	3%
hf-PGA score ≥3, N	319	124
Moderate (3)	80%	82%
Severe (4)	20%	18%
mNAPSI score >0, N	557	267
Mean mNAPSI score	19.8	17.2

^aPsO involving high-impact sites was not mutually exclusive. ^bAmong pts within each category of high-impact site PsO.


Baseline characteristics were generally similar between treatment groups within each high-impact site PsO cohort

Baseline Characteristics for Each Cohort of Pts With at Least Moderate High-Impact Site PsO	Scalp PsO: ss-IGA score ≥ 3		Genital PsO: sPGA-G score ≥ 3		Hand/foot PsO: hf-PGA score ≥ 3	
	ICO (N=935)	PBO (N=408)	ICO (N=334)	PBO (N=139)	ICO (N=319)	PBO (N=124)
Demographics						
Age, yrs	43.6 (14.9)	44.1 (14.6)	43.3 (13.8)	44.2 (15.0)	47.3 (14.2)	48.0 (13.6)
Female	318 (34%)	142 (35%)	103 (31%)	49 (35%)	100 (31%)	38 (31%)
Race, Asian / Black / White	21% / 2% / 75%	23% / 1% / 75%	19% / 1% / 77%	17% / 0% / 81%	22% / 2% / 76%	22% / 3% / 72%
BMI, kg/m ²	29.3 (6.5) ^a	29.3 (7.4) ^a	29.4 (6.6) ^b	29.6 (7.6) ^b	29.7 (6.6) ^c	30.2 (7.4) ^c
Disease Characteristics						
PsO duration, yrs	16.7 (12.8)	16.8 (12.7)	16.8 (13.0)	16.2 (11.3)	16.7 (12.1)	16.4 (13.1)
% of BSA with PsO	23.9 (14.9)	23.4 (15.0)	24.8 (14.7)	24.1 (15.2)	27.4 (17.7)	27.3 (17.5)
IGA score						
Moderate (3)/Severe (4)	74% / 25%	73% / 26%	73% / 26%	71% / 28%	69% / 31%	70% / 30%
PASI (0-72)	19.3 (7.7)	19.1 (7.9)	19.6 (7.6)	19.8 (7.9)	20.7 (8.9)	20.4 (9.4)
Prior PsO Treatments						
Phototherapy (PUVA and UVB)	34%	31%	35%	27%	33%	28%
Systemic therapy ^d	73%	71%	72%	75%	67%	68%
Biologic therapy ^e	30%	33%	31%	33%	26%	40%

Data shown are mean (SD), unless otherwise noted. ^aICO N=930/PBO N=405. ^bICO N=333/PBO N=138. ^cICO N=318/PBO N=123. ^dConventional nonbiologic systemics, novel nonbiologic systemics, 1,25-vitamin D3 and analogues, phototherapy, and biologics. ^eAdalimumab, alefacept, briakinumab, brodalumab, certolizumab pegol, efalizumab, etanercept, guselkumab, infliximab, ixekizumab, natalizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab. **BMI**=body mass index, **PUVA**=psoralen plus ultraviolet A, **SD**=standard deviation, **UVB**=ultraviolet B.

Scalp PsO: ICO demonstrated high rates of clearance through W24


- After transitioning to ICO at W16, PBO-randomized pts demonstrated consistent patterns of response through W24



^aAmong pts with a baseline ss-IGA score ≥ 3 . ^bNonresponder imputation. Confidence intervals based on the normal approximation confidence limits. CI=confidence interval.

Genital PsO: ICO demonstrated high rates of clearance through W24


- After transitioning to ICO at W16, PBO-randomized pts demonstrated consistent patterns of response through W24



^aAmong pts with a baseline sPGA-G score ≥ 3 . ^bNonresponder imputation. Confidence intervals based on the normal approximation confidence limits.

Hand/foot PsO: ICO demonstrated high rates of clearance through W24


- After transitioning to ICO at W16, PBO-randomized pts demonstrated consistent patterns of response through W24



^aAmong pts with a baseline hf-PGA score ≥ 3 . ^bNonresponder imputation. Confidence intervals based on the normal approximation confidence limits.

Nail PsO: ICO-randomized pts exhibited increasing mean percent mNAPSI improvement through W24

- PBO-randomized pts demonstrated substantial improvement in nail PsO within 8 weeks of starting ICO



^aAmong pts with a baseline mNAPSI score >0.

The ICO AE profile was similar to PBO through W16 in pts with at least moderate high-impact site PsO

Pooled Safety in Pts With at Least Moderate High-Impact Site PsO Through W16^a	ICO (N=1068)	PBO (N=460)
Mean weeks of follow-up	15.9	15.6
Any AE	516 (48%)	233 (51%)
Serious AE	15 (1%)	9 (2%)
AE leading to discontinuation	21 (2%)	15 (3%)
Infection	252 (24%)	114 (25%)
Serious infection	0%	1 (<1%)
Malignancy	5 (<1%)	1 (<1%)

Key Takeaways



In large pooled phase 3 cohorts of adults and adolescents with at least moderate high-impact site PsO and/or nail PsO, the targeted oral peptide ICO demonstrated:

✓ **High rates of scalp, genital, and hand/foot PsO clearance at W24**

✓ **72-84% achieved clear/almost clear high-impact site PsO**

✓ **57-73% achieved completely clear high-impact site PsO**

✓ **Substantial mean mNAPSI improvement (50%) at W24**



Consistent with findings from each study, the ICO AE profile was similar to PBO through W16⁴⁻⁶

References

1. Blauvelt A. *J Psoriasis Psoriatic Arthritis*. 2023;8:100-6.
2. Strober B. *J Am Acad Dermatol*. 2020;82:117-22.
3. Fourie AM. *Sci Rep*. 2024;14:17515.
4. Bissonnette R. *N Engl J Med*. 2025;393:1784-95.
5. Stein Gold L. *Lancet*. 2025;406:1363-74.
6. Gooderham M. *NEJM Evid*. 2025;4:EVIDoa2500155. doi: 10.1056/EVIDoa2500155.
7. Soung J. Presented at: AAD Innovation Academy; July 12, 2025; Chicago, IL, USA.

Acknowledgments

- > Medical writing support was provided by R. Contento, PharmD of Johnson & Johnson under the direction of the authors in accordance with Good Publication Practice guidelines (*Ann Intern Med.* 2022;175:1298-1304).
- > This presentation was sponsored by Johnson & Johnson.