

Structural Damage Progression and Clinical Response to Guselkumab in Participants With Active and Erosive Psoriatic Arthritis: Post-hoc Analysis at Week 24 of the Phase 3b, Randomized, Double-Blind, Placebo-Controlled APEX Study

Christopher T. Ritchlin,¹ Proton Rahman,² Laura C. Coates,³ Alexa P. Kollmeier,⁴ Yusang Jiang,⁵ Luis Calderon,⁶ Soumya D. Chakravarty,^{6,7} Elizabeth Adamson,⁶ Karissa Lozenski,⁶ Arthur Kavanaugh,⁸ Philip J. Mease^{9,10}

¹Department of Medicine, Allergy/Immunology and Rheumatology, University of Rochester Medical Center, Rochester, NY, USA; ²Faculty of Medicine, Division of Rheumatology, Memorial University of Newfoundland, St. Johns, NL, Canada; ³Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Botnar Research Centre, Oxford, UK; ⁴Johnson & Johnson, San Diego, CA, USA; ⁵Johnson & Johnson, Spring House, PA, USA; ⁶Johnson & Johnson, Horsham, PA, USA; ⁷Drexel University College of Medicine, Philadelphia, PA, USA; ⁸Center for Innovative Therapy, Division of Rheumatology, Allergy, Immunology, University of California San Diego, La Jolla, CA, USA; ⁹Rheumatology Research, Providence Swedish Medical Center, Seattle, WA, USA; ¹⁰University of Washington School of Medicine, Seattle, WA, USA



Scan the QR code. The QR code is intended to provide scientific information for individual reference, and the information should not be altered or reproduced in any way.

Background


Guselkumab (GUS) inhibition of structural damage progression in psoriatic arthritis (PsA)

- Uncontrolled inflammation in PsA increases the risk of structural joint damage, which is associated with long-term disability and greater impairments in physical function and health-related quality of life
- GUS is a fully human, dual-acting,¹ monoclonal antibody that selectively inhibits the interleukin (IL)-23p19 subunit
- In the phase 3b APEX study, GUS demonstrated efficacy across PsA domains and significantly inhibited structural damage progression compared with placebo (PBO) in participants (pts) with active and erosive PsA at Week (W) 24²

Objective

Evaluate the relationship between clinical response status and structural damage progression at W24 among GUS-randomized pts in APEX

APEX Study Design and Methods



ACR20/50/70=≥20%/≥50%/≥70% improvement in the American College of Rheumatology criteria; CASPAR=CASification criteria for Psoriatic Arthritis; cDAPSA=clinical Disease Activity Index for Psoriatic Arthritis; f/u=follow-up; GUS=guselkumab; HAQ-DI=Health Assessment Questionnaire-Disability Index; LDA=low disease activity; LSM=least squares mean; MD=minor disruption (Ukraine/Russia/Cuba); MDA=minimal disease activity; MI=multiple imputation; ND=neutral disorder (COVID-19 site access restrictions); NSAIDs=nonsteroidal anti-inflammatory drug; PBO=placebo; PsA=psoriatic arthritis; PsO=psoriasis; Pts=participants; Q4W=every 4 weeks; Q8W=every 8 weeks; R=randomization; REM=remission; SJC=swollen joint count; TJC=tender joint count; VAS=visual analog scale; vdH-S=van der Heide-Sharp; W=week.


Results

Baseline characteristics were well balanced across treatment groups

	GUS Q4W (N=273)	GUS Q8W (N=371)	PBO (N=376)	Total (N=1020)
Demographics				
Age, years	52.2 (13.2)	53.2 (12.9)	53.5 (13.0)	53.0 (13.0)
Male	55%	54%	57%	55%
Weight, kg	85.6 (20.1)	83.2 (17.4)	83.1 (18.2)	83.8 (18.5)
BMI, kg/m ²	29.4 (6.0)	29.0 (5.6)	28.9 (5.7)	29.1 (5.7)
PsA Characteristics				
PsA Disease Duration, years	7.5 (7.1)	7.2 (7.6)	7.2 (6.9)	7.3 (7.2)
SJC [0–66]	11.6 (9.4)	12.1 (8.5)	11.8 (8.9)	11.9 (8.9)
TJC [0–68]	21.2 (14.6)	20.6 (13.4)	20.5 (13.9)	20.7 (13.9)
Pt-reported pain (VAS; 0–100 mm)	59.2 (22.2)	59.3 (21.0)	59.4 (20.6)	59.3 (21.1)
HAQ-DI [0–3]	1.2 (0.7)	1.2 (0.6)	1.2 (0.7)	1.2 (0.7)
DAPSA	46.4 (24.3)	46.1 (22.1)	46.0 (22.8)	46.1 (22.9)
cDAPSA	44.7 (23.5)	44.6 (21.6)	44.3 (22.3)	44.5 (22.4)
CRP, mg/dL	1.7 (2.9)	1.5 (2.0)	1.7 (2.5)	1.6 (2.5)
Enthesitis / Dactylitis	58% / 44%	59% / 39%	59% / 45%	58% / 43%
Mean LEI [1–6]/DSS [1–60]	3.2 / 10.8	3.0 / 11.0	3.0 / 10.2	3.1 / 10.6
Radiographic Characteristics				
PsA-modified vdH-S score [0–528]	27.7 (47.6)	26.7 (43.4)	26.8 (42.2)	27.0 (44.1)
Erosion score [0–320]	13.7 (24.3)	13.4 (21.9)	13.4 (20.7)	13.5 (22.1)
JSN score [0–208]	14.0 (24.2)	13.3 (22.8)	13.4 (22.4)	13.5 (23.0)


Values are reported as mean (SD) for pts with nonmissing data unless otherwise noted. BMI=body mass index; cDAPSA=clinical Disease Activity Index for Psoriatic Arthritis; CRP=C-reactive protein; DAPSA=Disease Activity Index for Psoriatic Arthritis; DSS=Disability Severity Score; GUS=guselkumab; HAQ-DI=Health Assessment Questionnaire-Disability Index; JSN=joint space narrowing; LEI=Leeds Enthesitis Index; PBO=placebo; PsA=psoriatic arthritis; Pts=participants; Q4W=every 4 weeks; Q8W=every 8 weeks; SD=standard deviation; SJC=swollen joint count; TJC=tender joint count; VAS=visual analog scale; vdH-S=van der Heide-Sharp; W=week.

At W24, GUS-randomized pts who achieved response in joint efficacy measures had numerically lower levels of structural damage progression




ACR20/50/70=≥20%/≥50%/≥70% improvement in the American College of Rheumatology criteria; LSM=least squares mean; PsA=psoriatic arthritis; Q4W=every 4 weeks; Q8W=every 8 weeks; vdH-S=van der Heide-Sharp; W=week.

At W24, GUS-randomized pts who achieved low or minimal disease activity by composite indices had numerically lower levels of structural damage progression



cDAPSA=clinical Disease Activity Index for Psoriatic Arthritis; DAPSA=Disease Activity Index for Psoriatic Arthritis; LDA=low disease activity; LSM=least squares mean; MDA=minimal disease activity; PsA=psoriatic arthritis; Q4W=every 4 weeks; Q8W=every 8 weeks; vdH-S=van der Heide-Sharp; W=week.

At Week 24, GUS-randomized pts who achieved normalized physical function and minimal pain had numerically lower levels of structural damage progression



^aAmong pts who had HAQ-DI score <0.5 at baseline. ^bAmong pts who had pt-reported pain VAS <15 at baseline. HAQ-DI=Health Assessment Questionnaire-Disability Index; LSM=least squares mean; PsA=psoriatic arthritis; Pts=participants; Q4W=every 4 weeks; Q8W=every 8 weeks; VAS=visual analog scale; vdH-S=van der Heide-Sharp; W=week.