Long-Term Adjunctive Lumateperone Treatment in Major Depressive Disorder: Results From a Six-Month Open-Label Extension Study

Willie R. Earley, MD; Suresh Durgam, MD; Susan G. Kozauer, MD; Changzheng Chen, PhD

Intra-Cellular Therapies, Inc., a Johnson & Johnson Company, Bedminster, NJ, USA

Intra-Cellular Therrander

ABSTRACT

Background:

Current treatments for major depressive disorder (MDD) are limited by side effects including weight gain and metabolic disturbances. Approximately 50% of patients with MDD have inadequate response to antidepressant therapy (ADT) leading to reduced adherence, functioning, and quality of life.

Lumateperone is an FDA-approved, atypical antipsychotic to treat schizophrenia and depressive episodes associated with bipolar I or II disorder. The efficacy and safety of lumateperone 42mg adjunctive to ADT in patients with MDD with inadequate ADT response was recently demonstrated in 2 Phase 3 trials (Study 501 NCT04985942; Study 502 NCT05061706). This Phase 3 open-label extension trial, Study 503 (NCT05061719), investigated the long-term safety of adjunctive lumateperone 42mg in patients who completed Study 501 or 502.

Methods:

Study 501 and 502 enrolled patients (18-65y) meeting DSM-5 criteria for MDD with inadequate response to 1-2 adequate courses of ADT in the current depressive episode and Montgomery-Åsberg Depression Rating Scale (MADRS) Total score ≥24 and Clinical Global Impression Scale-Severity (CGI-S) score ≥4. Patients who safely completed the 6-week double-blind treatment period could enroll in Study 503 to receive 26-week, open-label, oral, once-daily lumateperone 42mg adjunctive to continued ADT. The primary endpoint was safety and tolerability of lumateperone 42mg, measured by adverse events (AEs), extrapyramidal symptoms (EPS), suicidality, and changes in laboratory parameters, vital signs, and electrocardiogram (ECG) measures. The secondary endpoint was improvement/maintenance of depressive symptoms, measured by MADRS Total score and CGI-S score change from Study 501 or 502 baseline to Week 26 of open-label treatment.

Results:

Of 809 patients enrolled and treated, 684 (84.5%) completed the treatment period. Treatment-emergent AEs (TEAEs) occurred in 548 patients (67.7%), with 292 (36.1%) experiencing a drug-related TEAE. Treatment discontinuation due to an AE occurred in 7.4% of patients. TEAEs reported in ≥5% of patients were headache (16.6%), dizziness (10.6%), dry mouth (8.0%), nausea (7.7%), somnolence (7.2%), diarrhea (6.2%), and nasopharyngitis (5.2%). Most TEAEs (98.9%) reported were mild to moderate in severity. The rates of EPS-related TEAEs based on broad standard Medical Dictionary for Regulatory Activities query were low (3.8%) with no increase in EPS scales. Mean changes from baseline to end of treatment were minimal for body morphology, cardiometabolic laboratory values, prolactin levels, pulse rate, blood pressure, and ECG measures. No patients reported emergence of serious suicidal ideation or suicidal behavior during the study. Symptoms of depression improved as measured by mean change from baseline to Week 26 in MADRS Total score (-22.9) and CGI-S score (-2.7).

Conclusions:

Lumateperone 42mg adjunctive to ADT was safe and effective during 26-week treatment in patients with MDD and inadequate ADT response.

The following information concerns a use that has not been approved by the U.S. Food and Drug Administration.

BACKGROUND

- Major depressive disorder (MDD) is a highly burdensome illness that is associated with functional impairment, psychiatric and medical comorbidities, and reduced quality of life¹
- MDD is often chronic, with high risk of relapse and recurrence that may require long-term or lifetime treatment²
- Most patients fail to achieve remission (≈75%) or response (≈60%) with first-line antidepressant (ADT) treatment and required treatment switch or adjunctive treatment³ - Inadequate ADT response is associated with increased hospitalization and suicide risk and greater impairments in functioning⁴
- Currently, the only FDA-approved adjunctive treatment options for MDD are atypical antipsychotics that have safety and tolerability
- concerns that impact short- and long-term medication adherence, including weight gain, cardiometabolic disturbances, and extrapyramidal symptoms (EPS)5
- Lumateperone is a mechanistically novel antipsychotic that is currently FDA-approved to treat schizophrenia and depressive episodes associated with bipolar I or bipolar II disorder as monotherapy and as adjunctive therapy with lithium or valproate^{6,7}
- Lumateperone is a simultaneous modulator of serotonin, dopamine, and glutamate neurotransmission⁶
- Specifically, lumateperone is a potent serotonin 5-HT₂₄ receptor antagonist, a dopamine D₂ receptor presynaptic partial agonist and postsynaptic antagonist, a D₁ receptor-dependent indirect modulator of AMPA and NMDA currents, and a serotonin reuptake inhibitor⁶ - This novel mechanism of action with multimodal effects may confer robust efficacy with improved safety and tolerability compared with current treatment options
- The efficacy and safety of lumateperone 42 mg adjunctive to ADT was recently demonstrated in 2 Phase 3 trials (Study 501 NCT04985942; Study 502 NCT05061706)8,9
- This Phase 3 open-label extension (OLE) trial, Study 503 (NCT05061719), examined the long-term safety and antidepressant effects of
- adjunctive lumateperone 42 mg in patients with MDD who had completed Study 501 or 502

METHODS

- Studies 501 and 502 enrolled patients aged 18-65 years, meeting DSM-5 criteria for MDD with inadequate response to 1-2 adequate courses of ADT in the current depressive episode and Montgomery-Åsberg Depression Rating Scale (MADRS) Total score ≥24 and Clinical Global Impression Scale-Severity (CGI-S) score ≥4
- Patients who safely completed the 6-week double-blind treatment period could enroll in Study 503, which was a long-term OLE study to support FDA approval; all patients received open-label, oral, once-daily lumateperone 42 mg adjunctive to continued ADT
- During Study 503, lumateperone 42 mg was administered once daily in the evening for 26 weeks
- The primary endpoint was safety and tolerability of lumateperone 42 mg, measured by adverse events (AEs), EPS, suicidality, and changes in laboratory parameters, vital signs, physical examinations, and electrocardiogram (ECG) measures
- EPS were assessed using a narrow standard MedDRA query for EPS-related treatment-emergent AEs (TEAEs) and the clinician-rated Abnormal Involuntary Movement Scale (AIMS), Barnes Akathisia Rating Scale (BARS), and Simpson-Angus Scale (SAS)
- The secondary endpoint was efficacy for depression symptoms, as measured by MADRS Total score and CGI-S score change from Study 501 or 502 baseline to Week 26 of open-label treatment

RESULTS

Patient Population

• Of the 809 patients enrolled in the OLE safety population, 684 patients (84.5%) completed the treatment period (**Table 1**) - The most common reasons for treatment discontinuation were adverse events (7.4%) and withdrawal of consent (5.1%)

Table 1. Patient Disposition of Safety Population

	Lumateperone 42 mg + ADT
Enrolled in OLE, n	812
Safety population, n	809
Discontinued treatment, n (%) ^a	125 (15.5)
Adverse event	60 (7.4)
Patient withdrew consent	41 (5.1)
Protocol violation	8 (1.0)
Lack of efficacy	7 (0.9)
Lost to follow-up	5 (0.6)
Other	4 (0.5)
Completed the OLE treatment period n(%) ^a	684 (84.5)
Duran antican af maticanta turata d	

ADT, antidepressant therapy: OLE, open-label extension

- Demographics and baseline characteristics were similar to patients in the 6-week double-blind placebo-controlled treatment periods
- The most common selective serotonin reuptake inhibitor was citalopram/escitalopram (31.6%) and the most common serotoninnorepinephrine reuptake inhibitor was venlafaxine/desvenlafaxine (27.1%)
- Mean MADRS and CGI-S scores at double-bind baseline indicated moderate-to-severe depression (Table 2)

Table 2. Baseline Demographics and Disease Characteristics in Safety Population

	Lumateperone 42 mg + ADT (N=809)
Age, mean (range), years	46.2 (18-66)
Sex, n (%)	
Female	549 (67.9)
Male	260 (32.1)
Race, n (%)	
White	703 (86.9)
Asian	61 (7.5)
Black	37 (4.6)
Other	8 (1.0)
Hispanic or Latino ethnicity, n (%)	86 (10.6)
No. of lifetime depressive episodes, mean (range)	3.8 (1-36)
No. of lifetime of treatment failures n (%) ^a	
1	596 (73.7)
2	213 (26.3)
Background ADT during the OLE	
SSRI	535 (66.1)
SNRI	224 (27.7)
Other (bupropion)	50 (6.2)
MADRS Total score	
At DB baseline, mean (SD)	30.7 (3.8)
At OLE baseline, mean (SD)	18.2 (8.6)
CGI-S score	
At DB baseline, mean (SD)	4.7 (0.6)
At OLE baseline, mean (SD)	3.4 (1.1)

- Of 809 patients, 548 (67.7%) had ≥1 TEAE during the OLE (**Table 3**)
- Most TEAEs (>98%) were mild or moderate in severity Serious adverse events were 1.0%
- TEAEs that occurred in ≥5% of patients were headache, dizziness, dry mouth, nausea, somnolence, diarrhea, and nasopharyngitis
- TEAEs led to discontinuation in 7.4% of patients; only dizziness led to discontinuation in more than 1% of patients (9 patients, 1.1%)
- No suicidal behavior (per Columbia Suicide Severity Rating Scale) or suicidality serious AEs were reported in the study

Table 3. Adverse Events in Safety Population During OLE Period^a

Event, n (%)	Lumateperone 42 mg + ADT (N=809)
≥1 TEAE	548 (67.7)
Drug-related TEAE	292 (36.1)
SAE	8 (1.0)
Discontinued treatment due to AE	60 (7.4)
Deaths	0
TEAEs occurring in ≥5% of patients	
Headache	134 (16.6)
Dizziness	86 (10.6)
Dry mouth	65 (8.0)
Nausea	62 (7.7)
Somnolence	58 (7.2)
Diarrhea	50 (6.2)
Nasopharyngitis	42 (5.2)

ADT, antidepressant therapy; AE, adverse event; OLE, open-label extension; SAE, serious adverse event; TEAE, treatment-emergent adverse event.

Body Morphology, Metabolic, Prolactin, and Vital Sign Assessments

- Changes in body morphology were small (Table 4)
- Potentially clinically significant weight increase or decrease (≥7% change from baseline) was low and similar
- There were minimal changes in cardiometabolic parameters (**Table 4**)
- Mean changes in blood pressure, heart rate, and respiratory rate were minimal
- Mean changes in prolactin levels at end of treatment were low, similar to what was seen in double-blind studies, and not clinically relevant

Table 4. Mean Change in Body Morphology, Cardiometabolic Parameters, and Prolactin During OLE Period

	Lumate	Lumateperone 42 mg + ADT (N=809)	
	Baseline Mea	Mean Change From Baseline to EOT (S	
Veight, kg	78.96 (16.	.9) -0.16 (3.72)	
BMI, kg/m²	27.8 (5.06	6) -0.05 (1.33)	
Vaist circumference, cm	92.6 (13.7	7) -0.54 (5.50)	
Cholesterol, mg/dL			
Total	199.7 (42.1	10) -8.2 (32.30)	
LDL	138.4 (41.2	24) -9.6 (30.42)	
HDL	56.7 (16.9	0.1 (11.79)	
Triglycerides, mg/dL	137.3 (81.6	66) -0.2 (84.26)	
Glucose, mg/dL	93.3 (14.7)	75) 1.1 (15.56)	
nsulin, μIU/L	14.51 (19.9	96) -0.41 (22.41)	
Prolactin, ng/mL	10.07 (12.9	98) 1.13 (13.01)	
PCS criterion		n/N (%)	
≥7% increase in weight		66/779 (8.5)	
≥7% decrease in weight		75/779 (9.6)	

Extrapyramidal Symptoms Assessments

- There were no notable changes in EPS as assessed by clinician-rated scales during the study (Table 5)
- EPS as defined by categorical shifts in the BARS or SAS scales were rare (**Table 5**)
- The frequency of EPS-related TEAEs was 3.8%

Table 5. Changes in EPS-Related Scales During OLE Period


	Lumateperone 42 mg + ADT	
	Baseline Mean (SD)	Mean Change to EOT (SD)
BARS Total score	0.1 (0.52)	-0.1 (0.53)
AIMS Total score	0.0 (0.37)	-0.0 (0.38)
SAS Total score	0.1 (0.36)	-0.0 (0.38)
EPS defined by categorical shifts	n/N (%) ^a	
Parkinsonism: Baseline BARS ≤2 to BARS >2 during treatment	4/776 (0.5)	
Akathisia: Baseline SAS ≤3 to SAS >3 during treatment	14/770 (1.8)	

an = number of patients who met criteria at least once during the OLE period; N = the number of patients with available baseline assessments that did not meet the criteria and had at least one assessment during the OLE period. ADT, antidepressant therapy; AIMS, Abnormal Involuntary Movement Scale; BARS; Barnes Akathisia Rating Scale; EOT, end of treatment; EPS, extrapyramidal symptoms; OLE, open-label extension; SAS, Simpson-Angus Scale.

Assessment of Depression Symptoms

- Improvement in depression symptoms continued throughout the OLE period, as measured by change in MADRS Total score (Figure 1A) Similar improvement was seen in CGI-S score (data not shown)
- Most patients showed clinically meaningful improvements as assessed by MADRS response (79.5%) and remission (65.4%) criteria (Figure 1B)

Figure 1. Assessment of Depression Symptoms: Mean Change From Baseline in MADRS Total Score (A) and MADRS **Response and Remission Rates (B)**



ADT, antidepressant therapy; DB, double-blind; EOLTP, end of long-term treatment period; MADRS, Montgomery-Åsberg Depression Rating Scale; OLE, open-label extension

CONCLUSIONS

- Lumateperone 42 mg adjunctive to ADT was generally safe and well tolerated in long-term treatment in patients with MDD; there were no new safety findings, and AEs and safety parameters were consistent with the short-term 501 and 502 studies
- Over 26 weeks of treatment, lumateperone 42 mg adjunctive to ADT was associated with low risk of weight gain, cardiometabolic effects, and EPS
- In patients treated long-term with lumateperone 42 mg adjunctive to ADT, efficacy was maintained, and symptoms of depression improved throughout the study
- These results support the long-term safety and effectiveness of lumateperone 42 mg adjunctive to ADT in patients with MDD and inadequate ADT response

REFERENCES

- Proudman D, et al. *Pharmacoeconomics*. 2021;39(6):619-625. 7. Caplyta. Prescribing information. Intra-Cellular Therapies, 2. Lam RW, et al. *Can J Psychiatry*. 2024;69(9):641-687. 3. Pigott HE, et al. *BMJ Open*. 2023;13(7):e063095.
- 4. Knoth RL, et al. *Am J Manag Care*. 2010;16(8):e188-e196. 5. Spielmans GI, et al. *PLoS Med*. 2013;10(3):e100140310.
- 8. Durgam S, et al. "Lumateperone as Adjunctive Therapy ir 6. Titulaer J, et al. Eur Neuropsychopharmacol. 2022;62:22-35.
 - Patients With Major Depressive Disorder: Results From a Randomised, Double-blind, Phase 3 Trial," Poster presented at: European College of Neuropsychopharmacology (ECNP) Annual Congress, September 21-24, 2024, Milan, Italy.

Randomized, Double-Blind, Phase 3 Trial." Poster presented at: Psych Congress Annual Meeting, October 29-November

DISCLOSURES AND ACKNOWLEDGMENTS

WR Earley, S Durgam, SG Kozauer, and C Chen are full-time employees of Intra-Cellular Therapies, Inc., a Johnson & The authors thank all study investigators, research staff, and patients for their participation. The authors would like to thank Andew Cutler, MD, for providing guidance and consultation into the clinical interpretation of the data. Medical writing support was provided by Adam Ruth, PhD, of Nucleus Global, an Inizio company, funded by Intra-Cellular Therapies, Inc., a Johnson & Johnson company.

Scan the QR code for the full digital poster. The QR code is intended to provide scientific information for individual reference, and the information should not be altered

