

Efficacy of Adjunctive Lumateperone 42 mg Treatment Across a Broad Range of Depression Symptoms in Patients with Major Depressive Disorder

Cristoph U. Correll, MD¹⁻⁶; Suresh Durgam, MD⁷; Willie R. Earley, MD⁸; Ibrahim Turkoz, PhD⁹; John J. Sheehan, PhD, MBA, RPh¹⁰; Roger S. McIntyre, MD¹¹⁻¹²

¹Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Department of Psychiatry and Molecular Medicine, Hempstead, NY, USA; ²Northwell, New Hyde Park, NY, USA; ³Charité Universitätsmedizin Berlin, Department of Child and Adolescent Psychiatry, Berlin, Germany; ⁴German Center for Mental Health (DZPG), partner site Berlin, Berlin, Germany; ⁵Einsteiner Center for Population Diversity (ECPD), Berlin, Germany; ⁶Einsteiner Center for Youth Mental Health (ECYM), Berlin, Germany; ⁷Former employee of Intra-Cellular Therapies, a Johnson & Johnson company, Bedminster, NJ, USA; ⁸Intra-Cellular Therapies, a Johnson & Johnson company, Bedminster, NJ, USA; ⁹Johnson & Johnson, Titusville, NJ, USA; ¹⁰Johnson & Johnson Innovative Medicine, Raritan, NJ, USA; ¹¹Department of Psychiatry, University of Toronto, Toronto, ON, Canada; ¹²Department of Pharmacology and Toxicology, University of Toronto, Toronto, ON, Canada

Background

- Current treatments for major depressive disorder (MDD) may not resolve all symptoms, and the presence of residual symptoms predicts disease relapse and recurrence as well as reduced quality of life and functional impairment^{1,3}
- Lumateperone is an atypical antipsychotic indicated in adults for: (1) treatment of MDD as adjunct to antidepressant therapy (ADT); (2) treatment of depressive episodes associated with bipolar I or II disorder (bipolar depression), as monotherapy and as adjunctive therapy with lithium or valproate; (3) and treatment of schizophrenia⁴
- Lumateperone simultaneously modulates serotonin, dopamine, and indirectly through D₂-receptor modulation, glutamatergic NMDA and AMPA receptors⁵
- Although the mechanism of action is unknown, lumateperone is a potent serotonin 5-HT_{2A} receptor antagonist, a dopamine D₂ receptor presynaptic partial agonist and postsynaptic antagonist, a D₁ receptor-dependent indirect modulator of glutamatergic AMPA and NMDA currents, and a serotonin reuptake inhibitor^{4,5}
- This novel mechanism of action with multimodal effects may confer robust efficacy with improved safety and tolerability
- The two pivotal Phase 3, randomised, double-blind, placebo-controlled, multicentre trials (Study 501, NCT4985942; Study 502, NCT05061706) demonstrated that lumateperone 42 mg/day + ADT significantly improved depressive symptoms compared with placebo + ADT, as measured by both clinician-rated and patient-rated outcomes, in patients with MDD with inadequate ADT response^{6,7}
- Lumateperone 42 mg/day + ADT had a favourable safety profile, with no notable changes in cardiometabolic parameters or body weight^{6,7}
- This pooled post hoc analysis of Studies 501 and 502 assessed the efficacy of lumateperone 42 mg + ADT on individual Montgomery-Åsberg Depression Rating Scale (MADRS) item scores to determine whether improvements in depressive symptoms were observed broadly across MADRS items or were driven by changes in only a limited number of symptoms


Methods

- Pooled data were analysed from Studies 501 and 502 involving adults (18-65 years) with DSM-5-defined MDD with inadequate response to 1 or 2 ADT in the current depressive episode, with a MADRS Total score ≥ 24 , Clinical Global Impression-Severity score ≥ 4 and Quick Inventory of Depressive Symptomatology-Self Report-16 item score ≥ 14
- Patients with a significant risk of suicidal behaviour were excluded
- Patients were randomised to 6 weeks of lumateperone 42 mg + ADT or placebo + ADT
- The change from baseline to Day 43 in MADRS Total score was analysed using a mixed-effects model for repeated measures in the modified intent-to-treat (mITT) population (all randomised patients who received ≥ 1 treatment dose, with a baseline and ≥ 1 postbaseline MADRS Total score)
- This post hoc analysis investigated change in individual MADRS items using analysis of covariance models in the mITT population
- Clinically meaningful change in individual MADRS items was defined as the proportion of patients with an improvement of ≥ 2 points from baseline
- The proportion of patients with MADRS item scores ≤ 1 (minimal or no symptoms) at Day 43 was also assessed

Results


- A total of 950 patients were included in the pooled mITT population (lumateperone 42 mg + ADT, n=471; placebo + ADT, n=479)
- Baseline characteristics were similar for the 2 groups
- The mean (range) age was 45.2 (18-65) years in the lumateperone 42 mg + ADT group and 45.8 (18-65) years in the placebo + ADT group
- Most patients were female (67.5% in the lumateperone 42 mg + ADT group and 67.8% in the placebo + ADT group) and White (85.8% in the lumateperone 42 mg + ADT group and 86.2% in the placebo + ADT group)
- At Day 43, lumateperone + ADT significantly improved MADRS Total score versus placebo + ADT (least squares mean difference [LSMD] vs placebo + ADT, -4.7 [95% CI, -5.8, -3.7]; effect size, -0.59; P<.0001)
- Mean baseline scores on individual MADRS items ranged from 4.1 (reported sadness; 'pervasive') to 0.4 (suicidal thoughts; 'enjoys life or takes it as it comes') in the lumateperone + ADT group (Figure 1)
- Median baseline scores on individual MADRS items ranged from 4.0 (reported sadness, apparent sadness, reduced sleep, concentration difficulties, lassitude, inability to feel) to 0 (suicidal thoughts) in the lumateperone + ADT group
- At Day 43, there were statistically significant improvements in all 10 MADRS items with lumateperone + ADT versus placebo + ADT (Figure 1)
- The largest improvement was in reduced sleep (LSMD, -0.90; 95% CI, -1.08, -0.71; P<.001)
- The smallest improvement was in suicidal thoughts (LSMD, -0.06; 95% CI, -0.12, 0.00; P=.042); this may be explained by the lower baseline values on this item stemming from the exclusion of patients with risk of suicidal behaviour
- Significant improvements with lumateperone + ADT versus placebo + ADT were seen from Day 8 for apparent sadness, reported sadness, inner tension, and reduced sleep and from Day 15 for lassitude and inability to feel; improvements were maintained through end of treatment (Figure 1)

Figure 1. Improvements in Individual MADRS Items From Baseline to Day 43




Negative change in score indicates improvement.
*P value is based on inferior significance (COST) P values with change from baseline as the dependent variable, treatment and study as factors, and baseline value as the covariate.
ANOVA, analysis of covariance; ADT, antidepressant therapy; LSMD, least squares mean difference; MADRS, Montgomery-Åsberg Depression Rating Scale.

- At Day 43, improvements of ≥ 2 points were observed across 9 of the 10 MADRS items in patients receiving lumateperone 42 mg + ADT (Figure 2)
- The exclusion of patients with risk of suicidal behaviour explains the low proportion of patients with ≥ 2 point improvement in suicidal thoughts



- At Day 43, the proportion of patients with absent or minimal individual symptoms increased across all items in the lumateperone 42 mg + ADT group vs the placebo + ADT group (Figure 3)
- The exclusion of patients with risk of suicidal behaviour explains the higher proportion of patients with a score of ≤ 1 on the suicidal thoughts item at Day 43

Figure 3. Proportion of Patients With a Score of ≤ 1 on Each MADRS Item at Day 43 With Lumateperone 42 mg + ADT vs Placebo + ADT



An item score of 0 indicates absent or minimal symptoms.
ADT, antidepressant therapy; MADRS, Montgomery-Åsberg Depression Rating Scale.

References:

1. Lai W, et al. *BMC Psychiatry*. 2025;25:310; 2. Nierenberg AA, et al. *Psychol Med*. 2010;40:41-50; 3. Whiston A, et al. *Front Psychiatry*. 2022;13:746678; 4. Caplyta (lumateperone). Prescribing information. Intra-Cellular Therapies, Inc; 2026; 5. Titulaer J, et al. *Eur Neuropsychopharmacol*. 2022;62:22-35; 6. Durgam S, et al. *J Clin Psychiatry*. 2025;86:2515848; 7. Durgam S, et al. *Am J Psychiatry*. 2025;182(12):1072-1082; 8. McIntyre RS, et al. *JAMA Psychiatry*. 2026;83(7):741-750.

Conclusions

- Lumateperone 42 mg + ADT demonstrated superior efficacy compared with placebo + ADT in improving a broad range of depression symptoms in patients with MDD with inadequate ADT response
- Early improvement from baseline was seen across key MADRS items with lumateperone 42 mg + ADT versus placebo + ADT and maintained through end of treatment
- More patients achieved ≥ 2 points improvement in individual MADRS items with lumateperone 42 mg + ADT versus placebo + ADT
- Lumateperone + ADT compared with placebo + ADT was associated with an increased percentage of patients with no or minimal symptoms as assessed using individual MADRS items, which accords with evidence of relatively large effect size improvement in overall depressive symptoms⁹

Acknowledgments

The authors thank all study investigators, research staff, and patients for their participation. Medical writing support was provided by Adam Ruth, PhD, of Nucleus Global, an Inizio company, funded by Intra-Cellular Therapies, a Johnson & Johnson company.

Disclosures

CU Correll has been a consultant and/or advisor to or has received honoraria from AbbVie, Alkermes, Allergan, Angelini, Arrow, Autopharm, Axsome, Boehringer Ingelheim, Bristol Myers Squibb, Cardio Diagnostics, Casasco, Cerevel, CNX Therapeutics, Cogstate, Compass Pathways, Dymatis, Dolor, Denovo, Draig, Eli Lilly, Eumetis Therapeutics, Gedeon Richter, GH Research, Hetero, Hikma, Holmusk, Intra-Cellular Therapies, Janssen/J&J, Karuna, LB Pharmaceuticals, Lundbeck, Mapi, MapLight, MedinCell, MedLink Global, Merck, Mindpax, Mitsubishi Tanabe Pharmaceuticals, Mylan, Neuma Therapeutics, Neuropharm, Neurocrine, Neurocris, Neurocris, NeuShen, Neusignal Therapeutics, Newron, Novo, Novo Nordisk, Orion Pharma, OM Pharma, Ouka, Otsuka, Phrenix, PPD Biotech, qCell, Recognify Life Sciences, Recordati, Relmada, Response Pharmaceuticals, Reviva, Roemmers, Rovi, Salada, Sanofi, Seqirus, Serina, Servier, Sunitomo Pharma America, Sunovion, Sun Pharma, Supernus, Symrise, Tabuk, Takeda, Teva, Terran, Tolmar, Vertex, Viatrix, and Xenon Pharmaceuticals. He provided expert testimony for Janssen, Lundbeck, Neurocris, and Otsuka. He served on a Data Safety Monitoring Board for Compass Pathways, Intra-Cellular Therapies, Relmada, Reviva, and Rovi. He has received grant support from Boehringer Ingelheim, Janssen, and Takeda. He received royalties from UpToDate and is also a stock option or stockholder in Cardio Diagnostics, Kuleon Bioscience, LB Pharmaceuticals, MedinCell, MedLink Global, Mindpax, Qantic, and Terran.

S Durgam is a former employee of Intra-Cellular Therapies, a Johnson & Johnson company.

WR Earley is a full-time employee of Intra-Cellular Therapies, a Johnson & Johnson company.

I Turkoz is an employee of Johnson & Johnson.

JJ Sheehan is a former employee of Johnson & Johnson Innovative Medicine.

RS McIntyre has received research grant support from CIBR/GACD/National Natural Science Foundation of China (NSFC) and the Milken Institute; and speaker/consultation fees from Lundbeck, Janssen, Alkermes, Neuma Therapeutics, Boehringer Ingelheim, Sage, Biogen, Mitsubishi Tanabe Pharma, Purdue, Pfizer, Otsuka, Takeda, Neurocris, Neurocris, Sunovion, Bausch Health, Axsome, Novo Nordisk, Kris Pharma, Sanofi, Eisai, Intra-Cellular Therapies, NewBridge Pharmaceuticals, Viatrix, AbbVie, and Atai Life Sciences.

Funding

This study was funded by Intra-Cellular Therapies, a Johnson & Johnson company.

Neuropsychiatry



Scan the QR code
The QR code is intended to provide scientific information for individual reference, and the information should not be altered or reproduced in any way.