

Lumateperone 42 mg in Major Depressive Disorder: Demographic and Clinical Subgroups Efficacy Analysis in a Phase 3 Randomized Placebo-Controlled Trial

Raffaele Migliore, MA¹; Suresh Durgam, MD²; Willie R. Earley, MD¹; Yifan Mo, PhD¹; Hassan Lakkis, PhD¹; Susan G. Kornstein, MD³

¹Intra-Cellular Therapies, a Johnson & Johnson company, Bedminster, NJ, USA; ²Former employee of Intra-Cellular Therapies, a Johnson & Johnson company, Bedminster, NJ, USA; ³Department of Psychiatry, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

Background

- Major depressive disorder (MDD) is a highly disabling, complex, and heterogeneous disease that burdens a diverse patient population^{1,3}
- In patients with MDD with inadequate response to antidepressant therapy (ADT), adjunctive atypical antipsychotic treatment is a recognised treatment strategy, warranting the need to evaluate its use across demographic and disease characteristics⁴
- Lumateperone is an atypical antipsychotic indicated for: the treatment of MDD in adults as adjunct to antidepressant therapy (ADT); the treatment of depressive episodes associated with bipolar I or II disorder (bipolar depression) in adults, as monotherapy and as adjunctive therapy with lithium or valproate⁵
- Lumateperone is a potent serotonin 5-HT_{2A} receptor antagonist, a dopamine D₂ receptor presynaptic partial agonist and postsynaptic antagonist, a D₁ receptor-dependent indirect modulator of glutamatergic AMPA and NMDA currents, and a serotonin reuptake inhibitor⁶
- This novel mechanism of action with multimodal effects may confer robust efficacy with improved tolerability compared with current treatment options
- The efficacy and safety of lumateperone 42 mg adjunctive to ADT were established in two Phase 3 randomised, double-blind, placebo-controlled trials (Study 501 [NCT04985942] and Study 502

[NCT05061706]) in patients with MDD with inadequate ADT response^{7,8}

- In Study 501, patients who received lumateperone 42 mg + ADT demonstrated consistent and clinically meaningful improvements in depression symptoms and disease severity compared with placebo + ADT across subgroups⁹
- In this post hoc analysis of Study 502, the efficacy of lumateperone 42 mg + ADT vs placebo + ADT was evaluated within various demographic and clinical subgroups of patients with MDD with inadequate ADT response

Methods

- Eligible patients (18-65 years) met DSM-5 criteria for MDD with inadequate response to 1-2 ADT courses in the current major depressive episode and had Montgomery-Åsberg Depression Rating Scale (MADRS) Total score ≥ 24 , Clinical Global Impression-Severity (CGI-S) score ≥ 4 , and Quick Inventory of Depressive

Symptomatology-Self Report-16 item (QIDS-SR-16) Total score ≥ 14

- Inadequate response to ADT was defined as <50% improvement with ≥ 6 weeks ADT monotherapy, as confirmed by the Antidepressant Treatment Response Questionnaire
- Patients were randomised 1:1 to 6-week oral lumateperone 42 mg or placebo adjunctive to ADT
- Efficacy was assessed in the overall population for the primary endpoint (change in MADRS Total score from baseline to Day 43) and key secondary endpoint (change in CGI-S score from baseline to Day 43) using a mixed-effects model for repeated measures (MMRM)
- The subgroup analysis of these endpoints was performed by demographic and disease characteristic subgroups using an MMRM (Table 1)
- Change in QIDS-SR-16 Total score from baseline to Day 43 was also assessed in the overall population and subgroups using an analysis of covariance

Table 1. Demographic and Baseline Disease Characteristic Subgroups

Demographic subgroups	Baseline disease characteristic subgroups
- Age: ≤ 40 or >40 years - Sex: male or female - Race: White or non-White - Ethnicity: Hispanic/Latino or not Hispanic/Latino - Region: US or non-US	- Disease severity: MADRS Total score <32 or ≥ 32 - Type of ADT: SSRI or SNRI/other - Number of ADT failures in the current episode: 1 or 2 - Presence/absence of DSM-5-defined anxious distress

ADT, antidepressant therapy; DSM-5, Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition; MADRS, Montgomery-Åsberg Depression Rating Scale; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor.

Results

Patient Population

- The modified intent-to-treat (mITT) population comprised 469 patients (lumateperone + ADT, n=232; placebo + ADT, n=237)
- Baseline demographics and clinical characteristics were similar between treatment groups
- The mean age of patients was 46 years
- The majority of patients were female (69.6%) and White (95.4%)

Efficacy

- The primary and key secondary endpoints were met, with lumateperone + ADT significantly improving MADRS Total score (least squares mean difference vs placebo [LSMD], -4.5; effect size [ES], -0.56; $P<.0001$) and CGI-S

score (LSMD, -0.5; ES, -0.51; $P<.0001$), respectively, at Day 43 vs placebo + ADT in the mITT population


- Lumateperone + ADT also significantly improved QIDS-SR-16 Total score at Day 43 in the intent-to-treat (ITT) population (LSMD, -2.2; ES, -0.45; $P<.0001$)

Subgroup Analysis

- Lumateperone + ADT significantly improved MADRS Total score from baseline to Day 43 compared with placebo + ADT across all demographic subgroups assessed except for race (White, $P<.0001$; non-White, $P=.8717$) and ethnicity (Hispanic/Latino, $P=.1468$; not Hispanic/Latino, $P<.0001$) (Figure 1A)
- Significant improvements in MADRS Total score from baseline to Day 43 were seen with lumateperone + ADT vs placebo + ADT across all baseline disease characteristics evaluated (Figure 1B)


- Significant improvement in CGI-S scores from baseline to Day 43 was consistently demonstrated with lumateperone + ADT vs placebo + ADT across all demographic and clinical subgroups, except for race (White, $P<.0001$; non-White, $P=.8376$) and ethnicity (Hispanic/Latino, $P=.1884$; not Hispanic/Latino, $P<.0001$) (Figures 2A and 2B)
- Lumateperone + ADT significantly improved QIDS-SR-16 Total scores from baseline to Day 43 in the age and sex demographic subgroups and across all baseline disease characteristic subgroups compared with placebo + ADT (Figures 3A and 3B)
- Significance was not reached for race (White, $P<.0001$; non-White, $P=.6556$), ethnicity (Hispanic or Latino, $P=.2585$; not Hispanic or Latino, $P<.0001$), or region (US, $P=.4518$; non-US, $P<.0001$)

Figure 1. Change From Baseline to Day 43 in MADRS Total Score Among (A) Demographic and (B) Disease Characteristic Subgroups (mITT Population)




* $P<.05$ ** $P<.01$ *** $P<.001$ **** $P<.0001$ MMRM in the mITT population.
n=Number of patients with Day 43 assessment; NI=Number of patients in each group at baseline.
ADT, antidepressant therapy; LSMD, least squares mean difference; MADRS, Montgomery-Åsberg Depression Rating Scale; mITT, modified intent-to-treat; MMRM, mixed-effects model for repeated measures; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor.

Figure 2. Change From Baseline to Day 43 in CGI-S Score Among (A) Demographic and (B) Disease Characteristic Subgroups (mITT Population)



* $P<.05$ ** $P<.01$ *** $P<.001$ **** $P<.0001$ MMRM in the mITT population.
n=Number of patients with Day 43 assessment; NI=Number of patients in each group at baseline.
ADT, antidepressant therapy; CGI-S, Clinical Global Impression-Severity; LSMD, least squares mean difference; MADRS, Montgomery-Åsberg Depression Rating Scale; mITT, modified intent-to-treat; MMRM, mixed-effects model for repeated measures; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor.

Figure 3. Change From Baseline to Day 43 in QIDS-SR-16 Total Score Among (A) Demographic and (B) Disease Characteristic Subgroups (ITT Population)



* $P<.05$ ** $P<.01$ *** $P<.001$ **** $P<.0001$ ANCOVA in the ITT population.
n=Number of patients with Day 43 assessment; NI=Number of patients in each group at baseline.
ADT, antidepressant therapy; ANCOVA, analysis of covariance; LSMD, least squares mean difference; MADRS, Montgomery-Åsberg Depression Rating Scale; ITT, intent-to-treat; QIDS-SR-16, Quick Inventory of Depressive Symptomatology-Self Report-16 item; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor.

References:

- Yan G, et al. *Psychiatry Res*. 2024;337:115958; 2. Su YA, Si T. *Gen Psychiatry*. 2022;35:e100724; 3. Liu Q, et al. *J Psychiatr Res*. 2020;126:134-140; 4. Rafeyan R, et al. *J Clin Psychiatry*. 2020;81:OT190378R3; 5. Caplaya (lumateperone), Prescribing Information. Intra-Cellular Therapies, Inc; 2026; 6. Titulaer J, et al. *Eur Neuropsychopharmacol*. 2022;42:22-35; 7. Durgam S, et al. *J Clin Psychiatry*. 2025;86:25mS848; 8. Durgam S, et al. *Am J Psychiatry*. 2025;182(12):1072-1082; 9. Durgam S, et al. Poster presented at the 64th Annual Meeting of the American College of Neuropsychopharmacology (ACNP), 12-15 January 2026, Nassau, Bahamas.

Conclusions

Lumateperone 42 mg adjunctive to ADT showed significant improvements over placebo adjunctive to ADT in depression symptoms and disease severity across most subgroups of patients with MDD enrolled in Study 502

These results further support adjunctive lumateperone 42 mg as an effective option for patients with MDD with inadequate ADT response, regardless of demographic or disease characteristics

Acknowledgments

The authors thank all study investigators, research staff, and patients for their participation. Medical writing support was provided by Joy Steenkamp, PhD, of Nucleus Global, an Inizio company, funded by Intra-Cellular Therapies, a Johnson & Johnson company.

Disclosures

R Migliore, WR Earley, Y Mo, and H Lakkis are full-time employees of Intra-Cellular Therapies, a Johnson & Johnson company.

S Durgam is a former employee of Intra-Cellular Therapies, a Johnson & Johnson company.

SG Kornstein receives research support from the National Institutes of Health and the National Science Foundation. She has served as a consultant for Lilly, Sage Therapeutics, Reunion Neuroscience, Reimada, AbbVie, Gerbera, and Arrivo BioVentures.

Funding

This study was funded by Intra-Cellular Therapies, a Johnson & Johnson company.

Previously presented at the American Psychiatric Association (APA) Annual Meeting, 16-20 May 2026, San Francisco, CA, USA.

Neuropsychiatry



Scan the QR code
The QR code is intended to provide scientific information for individual reference, and the information should not be altered or reproduced in any way.