

Age and frailty analyses of transplant-ineligible patients with newly diagnosed multiple myeloma in the phase 3 MAIA and CEPHEUS trials of daratumumab + lenalidomide-dexamethasone and bortezomib + lenalidomide-dexamethasone

Christopher P. Venner¹, Sonja Zweegman², Shaji K. Kumar³, Thierry Facon⁴, Aurore Perrot⁵, Philippe Moreau⁶, Noopur S. Raj⁷, Cyrille Hulin⁸, Supratik Basu⁹, Vania Hungria¹⁰, Yael C. Cohen¹¹, Hartmut Goldschmidt¹², George Wang¹³, Kasey Bolyard¹⁴, Lorena Lopez-Masi¹⁴, Matteo Loi¹⁵, Fredrik Borgsten¹⁴, Melissa Rowe¹⁶, Nizar J. Bahis¹⁷, Saad Z. Usmani¹⁸

¹Cross Cancer Institute, University of Alberta, Edmonton, and BC Cancer - Vancouver Centre, University of British Columbia, Canada; ²Amsterdam UMC, Vrije Universiteit Amsterdam, Cancer Center Amsterdam, The Netherlands; ³Mayo Clinic Rochester, MN, USA; ⁴University of Lille, CHU Lille, Service des Maladies du Sang, Lille, France; ⁵CHU de Toulouse, IUCT-O, Université de Toulouse, UPS, Service d'Hématologie, Toulouse, France; ⁶University Hospital Hôtel-Dieu, Nantes, France; ⁷Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA; ⁸Hôpital Haut Léveléque, University Hospital, Pessac, France; ⁹Royal Wolverhampton NHS Trust and University of Wolverhampton, CRN West Midlands, NIHR, Wolverhampton, UK; ¹⁰Clinica Médica São Germano, São Paulo, Brazil; ¹¹Tel Aviv Sourasky (Ichilov) Medical Center, Tel Aviv, Israel; ¹²GNMG-Study Group at University Hospital Heidelberg, Internal Medicine V, Heidelberg, Germany; ¹³Johnson & Johnson, Spring House, PA, USA; ¹⁴Johnson & Johnson, Raritan, NJ, USA; ¹⁵Johnson & Johnson, Leiden, The Netherlands; ¹⁶Johnson & Johnson, High Wycombe, UK; ¹⁷Arnie Charbonneau Cancer Research Institute, University of Calgary, Calgary, AB, Canada; ¹⁸Memorial Sloan Kettering Cancer Center, New York, NY, USA

Key Takeaway

Dara-based regimens improved efficacy outcomes across age and frailty subgroups in the MAIA and CEPHEUS trials, reinforcing Dara-based regimens as SOC in TIE NDMM regardless of age or frailty (based on the simplified frailty index). These data offer clinically relevant insights to help guide treatment selection for TIE patients

Conclusions

DRd or DVRd improved PFS, MRD-negativity ≥CR rates, and sustained MRD-negativity rates vs Rd or VRd arms across age and frailty subgroups in both trials

AEs were consistent with the known profile of each individual drug

In both trials, within treatment arms, there were trends toward increased incidence of some AEs in older and frail patients compared to younger or non-frail patients

Please scan QR code



https://www.congresshub.com/EHA2026/Oncology/Daratumumab/Usmani-Age
The QR code is intended to provide scientific information for individual reference and the information should not be altered or reproduced in any way.

Acknowledgments
The authors and Johnson & Johnson thank the patients who participated in this study, the staff members at the study sites, the data and safety monitoring committee, and the staff members involved in data collection and analyses. This study was funded by Johnson & Johnson. Medical writing support was provided by Erin Burns-Tidmore, PhD, of Eloquent Scientific Solutions, part of Emsion Spark, an Emsion Medical Communications agency, a part of Emsion Pharma Group, and funded by Johnson & Johnson. © 2026 American Society of Clinical Oncology, Inc. Reused with permission. This abstract was accepted and previously presented at the 2026 ASCO Annual Meeting. All rights reserved.

Disclosures
SZ Usmani reports Consulting/Advisory Role for AbbVie, Amgen, Bristol Myers Squibb, Celgene, Genetech, Gilead Sciences, GlaxoSmithKline, Johnson & Johnson, Karyopharm Therapeutics, Merck, and Takeda; Research Funding from Amgen, Array BioPharma, Bristol Myers Squibb, Celgene, GlaxoSmithKline, Johnson & Johnson, Merck, Pharmacosmetics, Sanofi, Seagen, and Skyline Diagnostics.

Introduction

- The MAIA (NCT02252172) and CEPHEUS (NCT03652064) trials demonstrated that daratumumab (Dara) combined with standard of care (SOC) bortezomib, lenalidomide, and dexamethasone (VRd) or lenalidomide and dexamethasone (Rd) significantly improved clinical outcomes versus SOC in transplant-ineligible (TIE) newly diagnosed multiple myeloma (NDMM)¹⁻³
- Frailty is a well-recognized, high-risk feature and predictor of survival outcomes in patients with multiple myeloma (MM)⁴
- Older patients with MM experience more complicated treatment choices and outcomes due to comorbidities and age-related physiological changes⁵
- In this subgroup analysis, we describe efficacy and safety in TIE patients from both trials by age and baseline frailty using the latest data cuts
 - MAIA: clinical cutoff, Oct 2021; median follow-up, 64.5 months
 - CEPHEUS: clinical cutoff, Oct 2025; median follow-up, 76.0 months

Methods

- MAIA and CEPHEUS were randomized, open-label, multicenter, phase 3 trials
- Patients were randomized 1:1 to receive DRd or Rd in MAIA and to receive DVRd or VRd in CEPHEUS
- CEPHEUS enrolled 395 patients with TIE or transplant-deferred NDMM. Only patients with TIE NDMM from CEPHEUS were included in this analysis
- Subgroups included:
 - Age (<70, 70-75, or ≥75 y)
 - Frailty (assessed using the IFM simplified frailty score at baseline)⁶
 - Nonfrail = score of 0/1
 - Frail = score ≥2
- Efficacy endpoints included progression-free survival (PFS), complete response or better (≥CR) rates, overall minimal residual disease (MRD) negativity (10⁻⁵ threshold), and sustained MRD negativity

Results

Patient Population

- MAIA¹ included 737 patients (DRd, n=368; Rd, n=369) with a median age of 73 y (range, 45-90)
 - 155 (21%) patients were <70 y, 261 (35%) 70-75 y, and 321 (44%) ≥75 y; 341 (46%) patients were frail, 88.0% of whom were ≥70 y
- CEPHEUS³ included 289 TIE patients (DVRd, n=144; VRd, n=145) with a median age of 72 y (range, 51-80); no patients were >80 y
 - 70 (24%) patients were <70 y, 133 (46%) 70-75 y, and 86 (30%) ≥75 y; 83 (29%) patients were frail, 86% of whom were ≥70 y
- Median treatment duration was up to 2 to 3 times longer in those receiving Dara-based regimens across age and frailty subgroups in both MAIA and CEPHEUS (Table 1)
 - Within treatment arms, treatment duration was longer in nonfrail patients, and also generally longer in younger patients

Results (cont)

Table 1: Median (range) duration of study treatment (months) in MAIA and CEPHEUS

	Overall		Frail		Nonfrail		<70 years		70-75 years		≥75 years	
MAIA												
	DRd (n=364)	Rd (n=365)	DRd (n=168)	Rd (n=166)	DRd (n=196)	Rd (n=199)	DRd (n=78)	Rd (n=76)	DRd (n=129)	Rd (n=130)	DRd (n=157)	Rd (n=159)
	47.5 (0.1-77.3)	22.6 (0.0-77.5)	39.1 (0.1-77.3)	20.7 (0.0-69.3)	54.6 (0.7-76.2)	25.8 (0.3-77.5)	59.5 (0.1-73.3)	21.0 (0.0-76.3)	49.2 (0.2-76.2)	25.9 (0.0-77.5)	40.8 (0.2-77.3)	20.1 (0.1-71.4)
CEPHEUS												
	DVRd (n=144)	VRd (n=142)	DVRd (n=48)	VRd (n=33)	DVRd (n=96)	VRd (n=109)	DVRd (n=35)	VRd (n=35)	DVRd (n=68)	VRd (n=65)	DVRd (n=41)	VRd (n=42)
	57.1 (0.1-81.9)	34.0 (0.5-81.0)	44.5 (0.5-80.7)	25.7 (0.5-79.4)	62.6 (0.1-81.9)	35.4 (0.5-81.0)	72.5 (3.6-78.8)	38.9 (0.5-81.0)	55.3 (0.1-81.9)	33.6 (1.2-79.9)	49.9 (2.1-80.7)	23.1 (0.5-78.9)

Efficacy

- DRd or DVRd consistently improved overall (Figure 1A, 2A) and sustained (Figure 1B, 2B) MRD-negativity in both MAIA (Figure 1) and CEPHEUS (Figure 2)
- Dara improved PFS in MAIA and CEPHEUS TIE patients across most age and frailty subgroups (Figure 3)

Figure 1: Overall and sustained MRD-negativity (10⁻⁵) ≥CR rates in MAIA




Figure 2: Overall and sustained MRD-negativity (10⁻⁵) ≥CR rates in CEPHEUS





Figure 3: PFS based on age and frailty in MAIA and CEPHEUS



References

- Facon T, et al. *N Engl J Med* 2019; 380: 2104-15.
- Facon T, et al. *Leukemia* 2025;39:942-50.
- Usmani SZ, et al. *Nature Med* 2025;31:1195-1202.
- Palumbo A, et al. *Blood* 2015;125(13):2068-74.
- Côté J, et al. *Front Oncol*. 2026;16:1632275.


Safety

- In MAIA, rates of grade 3/4 TEAEs were generally higher in frail vs nonfrail patients, and in older vs younger patients within treatment arms (Table 2). COVID-19 infections/pneumonia were reported in <2% of patients in the DRd arm across subgroups. There were no new safety signals beyond the typical neutropenia and infection TEAEs occurring with DRd
- In CEPHEUS, there were no apparent trends in the incidence of Grade 3/4 TEAEs (Table 2), including COVID-19, based on age or frailty. There were no new safety signals beyond the typical hematologic and infection TEAEs occurring with DVRd
- The incidence of TEAEs leading to study treatment discontinuation was slightly lower in those receiving Dara across age and frailty subgroups in MAIA and CEPHEUS (Table 2)
- The incidence of related serious TEAEs was similar in those receiving DRd or DVRd compared with Rd or VRd across most age and frailty subgroups in MAIA and CEPHEUS (Figure 4A-B)

Table 2: Most common grade 3/4 TEAEs of interest and treatment-related discontinuations in MAIA and CEPHEUS

	Frail		Nonfrail		<70 years		70-75 years		≥75 years	
MAIA										
	DRd (n=168)	Rd (n=166)	DRd (n=196)	Rd (n=199)	DRd (n=78)	Rd (n=76)	DRd (n=129)	Rd (n=130)	DRd (n=157)	Rd (n=159)
Any Grade 3/4 TEAE	161 (95.8)	152 (91.6)	188 (95.9)	172 (86.4)	72 (92.3)	62 (81.6)	127 (98.4)	111 (85.4)	150 (95.5)	151 (95.0)
Neutropenia	101 (60.1)	56 (33.7)	96 (49.0)	79 (39.7)	37 (47.4)	21 (27.6)	62 (48.1)	48 (36.9)	98 (62.4)	66 (41.5)
Infections & infestations	78 (46.4)	53 (31.9)	77 (39.3)	55 (27.6)	29 (37.2)	19 (25.0)	58 (45.0)	36 (27.7)	68 (43.3)	53 (33.3)
TEAE leading to study treatment discontinuation	27 (16.1)	40 (24.1)	26 (13.3)	47 (23.6)	9 (11.5)	11 (14.5)	20 (15.5)	32 (24.6)	24 (15.3)	44 (27.7)
CEPHEUS										
	DVRd (n=48)	VRd (n=33)	DVRd (n=96)	VRd (n=109)	DVRd (n=35)	VRd (n=35)	DVRd (n=68)	VRd (n=65)	DVRd (n=41)	VRd (n=42)
Any Grade 3/4 TEAE	48 (100)	29 (87.9)	87 (90.6)	97 (89.0)	32 (91.4)	30 (85.7)	62 (91.2)	58 (89.2)	41 (100)	38 (90.5)
Neutropenia	23 (47.9)	11 (33.3)	42 (43.8)	36 (33.0)	15 (42.9)	12 (34.3)	26 (38.2)	24 (36.9)	24 (58.5)	11 (26.2)
Infections & infestations	22 (45.8)	14 (42.4)	41 (42.7)	33 (30.3)	12 (34.3)	15 (42.9)	34 (50.0)	17 (26.2)	17 (41.5)	15 (35.7)
TEAE leading to study treatment discontinuation	5 (10.4)	7 (21.2)	9 (9.4)	26 (23.9)	2 (5.7)	7 (20.0)	7 (10.3)	12 (18.5)	5 (12.2)	14 (33.3)

Figure 4: Related serious TEAEs



- In MAIA, more patients died in the Rd vs DRd arm, regardless of frailty; more patients died of AEs in the frail vs nonfrail subgroup, regardless of treatment. Across age subgroups, more older patients died of progressive disease regardless of treatment (Table 3)
- In CEPHEUS, more frail patients died, regardless of treatment arm. Causes of death were similar between treatment arms in frail patients, whereas among nonfrail patients, more patients died of progressive disease in the VRd vs DVRd arm. No clear patterns emerged across age subgroups (Table 3)

Table 3: Causes of death in MAIA and CEPHEUS

	Frail		Nonfrail		<70 years		70-75 years		≥75 years	
MAIA										
	DRd (n=168)	Rd (n=166)	DRd (n=196)	Rd (n=199)	DRd (n=78)	Rd (n=76)	DRd (n=129)	Rd (n=130)	DRd (n=157)	Rd (n=159)
TEAE	22 (13.1)	21 (12.7)	7 (3.6)	10 (5.0)	7 (9.0)	3 (3.9)	6 (4.7)	10 (7.7)	16 (10.2)	18 (11.3)
Progressive disease	30 (17.9)	29 (17.5)	26 (13.3)	40 (20.1)	7 (9.0)	12 (15.8)	22 (17.1)	27 (20.8)	27 (17.2)	30 (18.9)
CEPHEUS										
	DVRd (n=48)	VRd (n=33)	DVRd (n=96)	VRd (n=109)	DVRd (n=35)	VRd (n=35)	DVRd (n=68)	VRd (n=65)	DVRd (n=41)	VRd (n=42)
TEAE	8 (16.7)	6 (18.2)	14 (14.6)	8 (7.3)	5 (14.3)	4 (11.4)	10 (14.7)	5 (7.7)	7 (17.1)	5 (11.9)
Progressive disease	6 (12.5)	4 (12.1)	2 (2.1)	13 (11.9)	2 (5.7)	4 (11.4)	2 (2.9)	9 (13.8)	4 (9.8)	4 (9.5)

