

# Daratumumab Plus Bortezomib, Lenalidomide, and Dexamethasone (DVRd) in Patients With Newly Diagnosed Multiple Myeloma: Final Analysis of Transplant-Ineligible (TIE) Patients in the Phase 3 CEPHEUS Study

**Saad Z. Usmani,<sup>1</sup> Thierry Facon,<sup>2</sup> Vania Hungria,<sup>3</sup> Nizar J. Bahlis,<sup>4</sup> Christopher P. Verner,<sup>5</sup> Marc Braunstein,<sup>6</sup> Ludek Pour,<sup>7</sup> Josep Marti,<sup>8</sup> Supratik Basu,<sup>9</sup> Yael C. Cohen,<sup>10</sup> Kenshi Suzuki,<sup>11</sup> Cyrille Hulin,<sup>12</sup> Sebastian Grosicki,<sup>13</sup> Meral Beksac,<sup>14</sup> Angelo Maiolino,<sup>15</sup> Jaclyn Stanziola,<sup>16</sup> Lorena Lopez-Masi,<sup>16</sup> Matteo Loi,<sup>17</sup> Melissa Rowe,<sup>18</sup> Sonja Zweegman<sup>19</sup>**

<sup>1</sup>Memorial Sloan Kettering Cancer Center, New York, NY, USA, <sup>2</sup>University of Lille, CHU Lille, Service des Maladies du Sang, Lille, France, <sup>3</sup>Clínica Médica São Germano, São Paulo, Brazil, <sup>4</sup>Arnie Charbonneau Cancer Research Institute, University of Calgary, Calgary, AB, Canada, <sup>5</sup>Cross Cancer Institute, University of Alberta, Edmonton, AB, Canada; BC Cancer–Vancouver Centre, University of British Columbia, Vancouver, BC, Canada, <sup>6</sup>Perlmutter Cancer Center, NYU Langone Health, New York, NY, USA, <sup>7</sup>University Hospital Brno, Brno, Czech Republic, <sup>8</sup>Hospital Universitario Mútua de Terrassa, Terrassa, Spain, <sup>9</sup>Royal Wolverhampton NHS Trust and University of Wolverhampton, CRN West Midlands, NIHR, Wolverhampton, UK, <sup>10</sup>Tel Aviv Sourasky (Ichilov) Medical Center, Tel Aviv, Israel; Gray Faculty of Medical and Health Sciences, Tel Aviv University, Tel Aviv, Israel, <sup>11</sup>Japanese Red Cross Medical Center, Tokyo, Japan, <sup>12</sup>Hôpital Haut Lévêque, University Hospital, Pessac, France, <sup>13</sup>School of Public Health, Medical University of Silesia, Katowice, Poland, <sup>14</sup>İstinye University, Ankara Liv Hospital, Ankara, Turkey, <sup>15</sup>Instituto Americas de Ensino, Pesquisa e Inovação, Rio de Janeiro, Brazil; Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil, <sup>16</sup>Johnson & Johnson, Raritan, NJ, USA, <sup>17</sup>Johnson & Johnson, Leiden, The Netherlands, <sup>18</sup>Johnson & Johnson, High Wycombe, UK, <sup>19</sup>Amsterdam UMC, Vrije Universiteit Amsterdam, Cancer Center Amsterdam, Amsterdam, The Netherlands

© 2026 American Society of Clinical Oncology, Inc. Reused with permission. This abstract was accepted and previously presented at the 2026 ASCO Annual Meeting. All rights reserved.

Presented by SZ Usmani at the European Hematology Association (EHA) Congress; June 11–14, 2026; Stockholm, Sweden

<https://www.congresshub.com/EHA2026/Oncology/Daratumumab/Usmani>

Scan the QR code

The QR code is intended to provide scientific information for individual reference, and the information should not be altered or reproduced in any way.



# CEPHEUS TIE Final Analysis: Key Takeaways

- After median follow-up of over 6 years (76.0 months), DVRd continued to show superiority vs VRd in all key endpoints, including:
  - Durable and deep responses, with 61.1% achieving overall MRD negativity at  $10^{-5}$  (vs. 40.0% VRd; OR 2.35; 95% CI 1.47–3.77) and a higher rate of sustained MRD negativity for  $\geq 12$  and  $\geq 24$  months
  - Nearly 60% of patients remained alive and progression-free at 72 months with DVRd, with median PFS not yet reached (vs. 50.2 months with VRd; HR 0.55; 95% CI 0.39–0.78)
  - No additional safety concerns compared with the ITT population in this older, frailer TIE subgroup

**The final CEPHEUS analysis results reinforce DVRd as the standard of care for patients with TIE NDMM, providing deeper and more durable responses than those receiving VRd**



# Introduction

- Daratumumab-based quadruplet therapy has transformed the NDMM treatment landscape and demonstrated improved survival outcomes for patients<sup>1-4</sup>
  - DVRd is a recommended treatment option for the treatment of TE<sup>5-7</sup> and TIE NDMM<sup>6,7</sup>
- The phase 3 CEPHEUS trial established DVRd as a new standard of care in TIE and TD NDMM<sup>4</sup>:
  - Improved depth and duration of response with higher rates of overall MRD ( $10^{-5}$ ) negativity and higher rates of sustained  $\geq 1$ -year and  $\geq 2$ -year MRD negativity
  - Improved PFS with a higher proportion of patients alive and progression-free at 4.5 years<sup>8</sup>
- **The majority of patients (~75%) enrolled in CEPHEUS were TIE – an older, frailer population vs ITT, with distinct clinical needs**

## TIE POPULATION at BL – AGE PROFILE (n=289)

**72 years**

Median age

**~76%**

Aged  $\geq 70$  years

**~30%**

Aged  $\geq 75$  years

**This final analysis of the CEPHEUS TIE subpopulation describes efficacy and safety outcomes of DVRd in TIE NDMM after a longer median follow-up of 76 months**

MAIA: NCT02252172; CEPHEUS: NCT03652064. BL, baseline; DRd, daratumumab, lenalidomide, and dexamethasone; DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; MRD, minimal residual disease; NDMM, newly diagnosed multiple myeloma; OS, overall survival; PFS, progression-free survival; Rd, lenalidomide and dexamethasone; TD, transplant deferred; TE, transplant eligible; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone.


1. Facon T, et al. *N Engl J Med* 2019;380:2104-15. 2. Sonneveld P, et al. *N Engl J Med* 2024;390:301-13. 3. Facon T, et al. *Leukemia* 2025;39:942-50. 4. Usmani, SZ, et al. *Nat Med* 2025;31:1195-202.

5. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Multiple Myeloma V.5.2026. © National Comprehensive Cancer Network, Inc. 2026. All rights reserved. Accessed March 24, 2026.

To view the most recent and complete version of the guideline, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way. 6. Rajkumar SV. *Am J Hematol* 2024;99:1802-24. 7. mSMART. Treatment for Multiple Myeloma v22. 8. Facon T, et al. Presented at ASCO 2025; Chicago, USA.



# CEPHEUS: Study Design




**This CEPHEUS final subgroup analysis includes only patients with TIE NDMM (~75% of the ITT population)**

<sup>a</sup>MRD was assessed via next-generation sequencing (clonoSEQ<sup>®</sup>; Adaptive Biotechnologies) using bone marrow aspirate samples obtained at baseline, at the time of suspected CR, and at 12, 18, 24, 30, and 36 months after the first dose and annually thereafter in patients with CR. CR, complete response; DRd, daratumumab, lenalidomide, and dexamethasone; DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; ECOG PS, Eastern Cooperative Oncology Group performance status; IMWG, International Myeloma Working Group; ITT, intent to treat; MRD, minimal residual disease; NDMM, newly diagnosed multiple myeloma; OS, overall survival; PFS, progression-free survival; Rd, lenalidomide and dexamethasone; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone. ClinicalTrials.gov Identifier: NCT03652064. Accessed April 7, 2025. Usmani, SZ, et al. *Nat Med* 2025;31:1195-202.



# CEPHEUS TIE Final Analysis: Overall and Sustained MRD-Negativity $\geq$ CR Rates

Overall MRD-negativity ( $\geq$ CR) rate<sup>a</sup>



At 10<sup>-5</sup>: OR 2.35; 95% CI 1.47–3.77; *P*=0.0004  
At 10<sup>-6</sup>: OR 2.27; 95% CI 1.39–3.71; *P*=0.0010

**DVRd demonstrated superior overall and sustained MRD negativity rates at 10<sup>-5</sup> and 10<sup>-6</sup> vs VRd**


The overall  $\geq$ CR rate was not changed from the primary analysis: 80.6% (DVRd) vs 61.4% (VRd) (OR 2.64; 95% CI 1.54–4.51; *P*=0.003)

<sup>a</sup>The proportion of patients who achieved MRD negativity and  $\geq$ CR. <sup>b</sup>Sustained MRD negativity was defined as 2 consecutive MRD negative reads  $\geq$ 12 months ( $\pm$ 1) or 24 months ( $\pm$ 3) apart with no MRD-positive result in between. CR, complete response; DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; MRD, minimal residual disease; OR, odds ratio; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone.




# CEPHEUS TIE Final Analysis: Overall and Sustained MRD-Negativity $\geq$ CR Rates

Overall MRD-negativity ( $\geq$ CR) rate<sup>a</sup>



At 10<sup>-5</sup>: OR 2.35; 95% CI 1.47–3.77; P=0.0004  
 At 10<sup>-6</sup>: OR 2.27; 95% CI 1.39–3.71; P=0.0010

Sustained MRD-negativity ( $\geq$ CR) rate  $\geq$ 12 months<sup>b</sup>



At 10<sup>-5</sup>: OR 2.40; 95% CI 1.47–3.91; P=0.0005  
 At 10<sup>-6</sup>: OR 3.01; 95% CI 1.73–5.24; P<0.0001

**DVRd demonstrated superior overall and sustained MRD negativity rates at 10<sup>-5</sup> and 10<sup>-6</sup> vs VRd**


The overall  $\geq$ CR rate was not changed from the primary analysis: 80.6% (DVRd) vs 61.4% (VRd) (OR 2.64; 95% CI 1.54–4.51; P=0.003)

<sup>a</sup>The proportion of patients who achieved MRD negativity and  $\geq$ CR. <sup>b</sup>Sustained MRD negativity was defined as 2 consecutive MRD negative reads  $\geq$ 12 months ( $\pm$ 1) or 24 months ( $\pm$ 3) apart with no MRD-positive result in between. CR, complete response; DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; MRD, minimal residual disease; OR, odds ratio; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone.




# CEPHEUS TIE Final Analysis: Overall and Sustained MRD-Negativity $\geq$ CR Rates

Overall MRD-negativity ( $\geq$ CR) rate<sup>a</sup>




At 10<sup>-5</sup>: OR 2.35; 95% CI 1.47–3.77; P=0.0004  
 At 10<sup>-6</sup>: OR 2.27; 95% CI 1.39–3.71; P=0.0010

Sustained MRD-negativity ( $\geq$ CR) rate  $\geq$ 12 months<sup>b</sup>



At 10<sup>-5</sup>: OR 2.40; 95% CI 1.47–3.91; P=0.0005  
 At 10<sup>-6</sup>: OR 3.01; 95% CI 1.73–5.24; P<0.0001

Sustained MRD-negativity ( $\geq$ CR) rate  $\geq$ 24 months<sup>b</sup>



At 10<sup>-5</sup>: OR, 2.62 (95% CI, 1.58–4.36); P=0.0002  
 At 10<sup>-6</sup>: OR, 2.45 (95% CI, 1.38–4.37); P=0.0020


**DVRd demonstrated superior overall and sustained MRD negativity rates at 10<sup>-5</sup> and 10<sup>-6</sup> vs VRd**

The overall  $\geq$ CR rate was not changed from the primary analysis: 80.6% (DVRd) vs 61.4% (VRd) (OR 2.64; 95% CI 1.54–4.51; P=0.003)

<sup>a</sup>The proportion of patients who achieved MRD negativity and  $\geq$ CR. <sup>b</sup>Sustained MRD negativity was defined as 2 consecutive MRD negative reads  $\geq$ 12 months ( $\pm$ 1) or 24 months ( $\pm$ 3) apart with no MRD-positive result in between. CR, complete response; DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; MRD, minimal residual disease; OR, odds ratio; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone.





# CEPHEUS TIE Final Analysis: Progression-Free Survival



**DVRd significantly improved PFS with 45% reduction in risk of disease progression or death vs VRd**



# CEPHEUS TIE Final Analysis: Overall MRD Negativity ( $10^{-5}$ ) With $\geq$ CR and PFS in Prespecified Subgroups



Treatment effect was generally consistent across subgroups



# CEPHEUS TIE Final Analysis: Safety

## Safety overview

Event, n (%)	DVRd (n=144)	VRd (n=142)
<b>Any TEAE</b>	144 (100.0)	142 (100.0)
Grade 3 and 4	135 (93.8)	126 (88.7)
Grade 5 non-COVID-19	18 (12.5)	13 (9.2)
Grade 5 COVID-19 <sup>a</sup>	6 (4.2)	1 (0.7)
<b>Exposure adjusted grade 5 TEAE rate, patient-months<sup>b</sup></b>	0.30/100	0.26/100
<b>Any serious TEAE</b>	109 (75.7)	99 (69.7)
<b>TEAE leading to discontinuation of all study treatment</b>	14 (9.7)	33 (23.2)
<b>Second primary malignancies</b>	20 (13.9)	20 (14.1)

## Common (≥5%) grade 3 or 4 TEAEs of interest


Event, n (%)	DVRd (n=144)	VRd (n=142)
<b>Neutropenia</b>	65 (45.1)	47 (33.1)
<b>Thrombocytopenia</b>	44 (30.6)	33 (23.2)
<b>Anemia</b>	16 (11.1)	14 (9.9)
<b>Diarrhea</b>	20 (13.9)	15 (10.6)
<b>Fatigue</b>	13 (9.0)	15 (10.6)
<b>COVID-19<sup>c</sup></b>	14 (9.7)	5 (3.5)
<b>Pneumonia</b>	26 (18.1)	19 (13.4)
<b>Peripheral sensory neuropathy</b>	Any Grade: 86 (59.7) Grade 3/4: 14 (9.7)	Any Grade: 91 (64.1) Grade 3/4: 12 (8.5)

**DVRd showed no additional safety concerns with longer follow-up**

<sup>a</sup>Deaths on or within 30 days of treatment. <sup>b</sup>Exposure-adjusted incidence rate: number of subjects with event per 100 patient-months at risk. Patient-months at risk = sum of exposure time until first TEAE occurrence or end of treatment for subjects without the event. <sup>c</sup>Group term. DVRd, daratumumab, bortezomib, lenalidomide, and dexamethasone; ITT, intent to treat; TEAE, treatment-emergent adverse event; TIE, transplant ineligible; VRd, bortezomib, lenalidomide, and dexamethasone.




# CEPHEUS TIE Final Analysis: Overall Survival



Subjects at risk

VRd	145	135	128	120	115	111	106	103	98	94	93	90	84	14	0
DVRd	144	139	130	127	121	116	115	113	109	106	101	92	90	23	0



Subjects at risk

VRd	145	135	128	120	115	111	106	103	98	94	93	90	84	14	0
DVRd	144	139	130	127	121	116	115	113	109	106	101	92	90	23	0

- Leading cause of death was unrelated AEs:
  - 23 (16.0%) DVRd patients and 15 (10.6%) VRd patients
- Of the total deaths in the TIE population, 11 (3.8%) were COVID-19 related
  - 8/144 (5.6%) DVRd and 3/142 (2.1%) VRd
- Deaths from progressive disease were more than double in the VRd arm
  - 8 (5.6%) DVRd and 17 (12.0%) VRd

**OS favored DVRd, especially when censoring for death due to COVID-19 (HR 0.74)  
DVRd halved myeloma-related deaths versus VRd**



# CEPHEUS TIE Subgroup: Conclusions

- After a median follow-up of over 6 years, the final analysis of the CEPHEUS trial demonstrated that DVRd continues to show superior efficacy over VRd across key endpoints in TIE patients
- In this post hoc analysis, DVRd improved depth and duration of response:
  - Overall MRD negativity: 61.1% vs 40.0% at  $10^{-5}$ ; 46.5% vs 27.6% at  $10^{-6}$
  - 12-month sustained MRD negativity: 49.3% vs 29.0% at  $10^{-5}$ ; 37.5% vs 16.6% at  $10^{-6}$
  - 24-month sustained MRD negativity: 44.4% vs 23.4% at  $10^{-5}$ ; 30.6% vs 15.2% at  $10^{-6}$
- Risk of disease progression or death was 45% lower for DVRd vs VRd; HR, 0.55
- OS favored DVRd, especially when censoring for death due to COVID-19 (HR, 0.74)
- No additional safety concerns were identified with longer follow-up in this older, frailer TIE subgroup

**Results of this final CEPHEUS subgroup analysis reinforce  
DVRd as the standard of care for the treatment of TIE NDMM**



# CEPHEUS: Acknowledgments

- We thank the patients who participated in this study and their families/caregivers, staff members at the study sites, and the data and safety monitoring committee
- This study was funded by Johnson & Johnson



<https://www.congresshub.com/EHA2026/Oncology/Daratumumab/Usmani>

Scan the QR code

The QR code is intended to provide scientific information for individual reference, and the information should not be altered or reproduced in any way.

