

CARVYKTI® Batch/Lot No.:

Date of CARVYKTI® Infusion:

Hospital and Associated Clinics Name:

After Hours Phone Number:

Office Phone Number:

Name of CARVYKTI® Treating Oncologist:

**This Patient has received CARVYKTI®.**

**IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW: CARVYKTI® therapy can cause cytokine release syndrome (CRS) or neurological toxicities, which may be fatal or life-threatening. CRS may involve multiple organ systems. If your patient reports any signs or symptoms as referenced on this card, please contact the patient's oncologist immediately for further information.**

**FOR HEALTHCARE PROFESSIONALS**

# CARVYKTI® PATIENT WALLET CARD

Carry this card with you at all times. **SHOW THIS CARD** to any healthcare professional involved in your care and if you go to the emergency room.

**IMPORTANT SAFETY INFORMATION FOR PATIENTS  
RECEIVING TREATMENT WITH CARVYKTI®**



07/25 cp-262512v7

Johnson & Johnson



# FOR THE PATIENT

Call your healthcare professional or get emergency help right away if you recognize any of these symptoms:

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• Leg and arm weakness with paralysis</li><li>• Fever (100.4°F or 38°C or higher)</li><li>• Chills or shaking chills</li><li>• Difficulty breathing</li><li>• Fast or irregular heartbeat</li><li>• Very low blood pressure</li><li>• Dizziness or light-headedness</li><li>• Muscle or joint pain</li><li>• Confusion or disorientation</li><li>• Difficulty speaking, reading, or writing</li><li>• Changes in balance or coordination</li><li>• Difficulty moving muscles of face and eyes</li><li>• Facial numbness</li></ul> | <ul style="list-style-type: none"><li>• Tingling, numbness, and pain of hands and feet</li><li>• Slower movements</li><li>• Shuffling feet or small steps when walking</li><li>• Personality changes (e.g., less talkative, disinterest in activities)</li><li>• Smiling, frowning, or showing emotion less</li><li>• Difficulty performing simple tasks (e.g., getting dressed, feeding oneself)</li><li>• Memory loss or foggy</li><li>• Smaller handwriting</li><li>• Tremor</li></ul> |
|---|---|

**You should always ask your doctor about taking other medications while taking CARVYKTI®.**

**Important to remember:** Stay close to the location where you received your CARVYKTI® infusion for at least 2 weeks. If you have any of these symptoms call your doctor or seek emergency medical attention right away! These are not all of the possible side effects of CARVYKTI®. Tell your doctor if you have any side effect that bothers you or does not go away.