


DARZALEX® (daratumumab)

What are the possible side effects of DARZALEX?

DARZALEX may cause serious side effects, including¹:

Infusion-related reactions



- Infusion-related reactions are common with DARZALEX.
- Serious allergic reactions and systemic reactions due to release of certain substances by your body can lead to death.
- If you have infusion-related reactions, your healthcare provider may temporarily stop your infusion or may completely stop treatment with DARZALEX.
- Get medical help right away if you get any of the following symptoms:
 - Shortness of breath or trouble breathing
 - Dizziness or lightheadedness (hypotension)
 - Cough
 - Wheezing
 - Heart beating faster than usual
 - Low oxygen in the blood (hypoxia)
 - Throat tightness or irritation
 - Runny or stuffy nose
 - Headache
 - Itching
 - High blood pressure
 - Eye pain
 - Nausea
 - Vomiting
 - Fever
 - Chills
 - Chest discomfort
 - Blurred vision

Changes in blood tests




- DARZALEX can affect the results of blood tests to match your blood type.
- These changes can last for up to 6 months after your final dose of DARZALEX.
- Your healthcare provider will do blood tests to match your blood type before you start treatment with DARZALEX.
- Tell all of your healthcare providers that you are being treated with DARZALEX before receiving blood transfusions.

DARZALEX® (daratumumab)

What are the possible side effects of DARZALEX?

Infections



- DARZALEX can cause serious life-threatening infections that can lead to death.
- Tell your healthcare provider right away if you develop a fever, trouble breathing, cough, burning or pain when you urinate, or any other signs and symptoms of infection during treatment.

Decreases in blood cell counts



- DARZALEX can decrease white blood cell counts, which help fight infections, and blood cells called platelets, which help to clot blood.
- Decreases in blood cell counts are common with DARZALEX but can be severe.
- Your healthcare provider will check your blood cell counts during treatment with DARZALEX.
- Tell your healthcare provider if you develop fever or have signs of bruising or bleeding.

Most common side effects of DARZALEX include¹:

- Cold-like symptoms (upper respiratory infection)
- Diarrhea
- Constipation
- Decreased red blood cells
- Nerve damage causing tingling, numbness, or pain
- Tiredness
- Swollen hands, ankles, or feet
- Nausea
- Cough
- Fever
- Shortness of breath
- Feeling weak



- These are not all the possible side effects of DARZALEX.¹
- Call your doctor for medical advice about side effects.¹
- You may report side effects to the Food and Drug Administration (FDA) at 1-800-FDA-1088.¹

This information is provided ONLY in response to your request and is not intended as medical advice, to promote the use of our product, or suggest using it in any manner other than as described in the PRESCRIBING INFORMATION. All decisions regarding your medical treatment should be made with your healthcare professional(s). Please contact your healthcare team(s) to discuss the information in this response.

Please refer to the attached full PRESCRIBING INFORMATION and the PATIENT INFORMATION for DARZALEX.¹

1. DARZALEX (daratumumab) [Prescribing Information]. Horsham, PA: Janssen Biotech, Inc.; <https://www.injilabels.com/package-insert/product-monograph/prescribing-information/DARZALEX-pi.pdf>.