INLEXZO™ (gemcitabine intravesical system) How will I receive INLEXZO?



 INLEXZO will be inserted and removed by your healthcare provider (HCP).¹



INLEXZO is inserted into your bladder through a tube called a urinary catheter 1 time every 3 weeks for up to 6 months (8 doses) and then 1 time every 12 weeks for up to 18 months (6 doses).¹



 INLEXZO is removed after 3 weeks of being inserted (3-week indwelling period).¹



 Your HCP will decide how many INLEXZO treatments you will receive.¹



Your HCP may give you antibiotics before INLEXZO is inserted or removed.¹



It is very important that you go to all of your appointments. If you miss any appointments, call your HCP as soon as possible to schedule your appointment.¹

Before receiving INLEXZO1:



Do not empty your bladder right before your procedure to insert INLEXZO.

After receiving INLEXZO1:



 Drink about 6 to 7 cups (1500 mL) of fluids per day during treatment with INLEXZO to make sure you produce enough urine for the medicine to be released into the bladder.



You can urinate normally. There is no need to hold your urine.



Avoid contact between your skin and urine.



To urinate, **males and females** should sit on the toilet and flush after each use.



 Wash your hands with water and soap and wash your genital area with water after each time you urinate.



 Wash clothing soiled with urine right away and separately from other clothing.

After your HCP removes INLEXZO1:



• Avoid contact between your skin and urine for at least 24 hours after INLEXZO is removed.

INLEXZO™ (gemcitabine intravesical system) How will I receive INLEXZO?

What should I avoid after INLEXZO is inserted?

- Tell your health care provider (HCP) that you have INLEXZO before having a type of scan called Magnetic Resonance Imaging (MRI).¹
 - INLEXZO may be used under specific conditions.¹
 - o Your HCP will give you an MRI Safety Information Card.



- Keep the card in a safe place and show it to all of your HCPs.
- o The card contains important information in case you need to have an MRI.
- Your HCP will review the information on the MRI Safety Information Card and determine the conditions they can safely do an MRI scan while INLEXZO is in your bladder.

This information is provided ONLY in response to your request and is not intended as medical advice, to promote the use of our product, or suggest using it in any manner other than as described in the PRESCRIBING INFORMATION. All decisions regarding your medical treatment should be made with your healthcare professional(s). Please contact your healthcare professional(s) to discuss the information in this response.