

SPRAVATO® evidence and value summary: ESCAPE-TRD



Major depressive disorder (MDD) after initial treatment: Current gaps and limitations



One-third of patients with MDD do not achieve remission^{1 a,b}

No significant improvement in remission at Step 2 is observed between switching within a class or to a different class.



High direct costs and health resource utilization^{2c}

1.8-3.6x higher annual all-cause direct costs compared to non-TRD MDD and non-MDD patients.



Substantial impact on productivity and activities of daily living^{3d}

25% of patients with TRD are unemployed, and 21.5% are on long-term disability.

SPRAVATO®: A different mechanism of action indicated for two subgroups of MDD

Indication⁴
SPRAVATO® is a non-competitive N-methyl D-aspartate (NMDA) receptor antagonist indicated for the treatment of:

- TRD in adults, as monotherapy or in conjunction with an oral antidepressant
- Depressive symptoms in adults with MDD with acute suicidal ideation or behavior in conjunction with an oral antidepressant

SPRAVATO® is available only through the **SPRAVATO® REMS** restricted program because of the risks of SAEs. SPRAVATO® is intended for use only in certified healthcare treatment centers and under the direct observation of HCPs. Patients treated with SPRAVATO® require HCP monitoring for at least 2 hours. SPRAVATO® must never be dispensed directly to a patient for home use.⁴

Limitations⁴

- The effectiveness of SPRAVATO® in preventing suicide or reducing suicidal ideation or behavior has not been demonstrated.
- Use of SPRAVATO® does not preclude the need for hospitalization if clinically warranted, even if patients experience improvement after an initial dose of SPRAVATO®.
- SPRAVATO® is not approved as an anesthetic agent.



Boxed warning: Sedation, dissociation, abuse and misuse, suicidal thoughts and behaviors, respiratory depression.⁴

Please refer to the full prescribing information for a complete listing of all adverse events, including other serious adverse events.

SPRAVATO®: Clinical evidence: ESCAPE-TRD directly compares the efficacy of SPRAVATO® to quetiapine XR in patients with TRD receiving ongoing SSRI/SNRI⁵


ESCAPE-TRD: A 32-week, Phase 3b, randomized, open-label, rater-blinded comparative trial of SPRAVATO® vs. quetiapine XR in patients with TRD. The treatment period consisted of an 8-week acute phase followed by a 24-week maintenance phase.⁵

- ESCAPE-TRD was an international study in which SPRAVATO® was dosed according to its EMA summary of product characteristics. A post-hoc analysis evaluated a subgroup of patients with TRD who received treatment in accordance with US prescribing information (patients aged 18-64 years who received flexibly dosed SPRAVATO® 56 or 84 mg, consistent with US label dosing).⁶



In a cost-per-remitter analysis based on clinical outcomes from ESCAPE-TRD in which non-responders initiated rTMS or APS augmentation as alternative treatments in equal proportions.⁹

In a post-hoc subgroup analysis of patients in ESCAPE-TRD who received treatment with SPRAVATO® in accordance with US Prescribing Information⁹ compared to QUE XR^f, WPL in both treatment arms was assessed from baseline to Week 32. Annual cost-savings were extrapolated from results.¹⁰



^aThe STAR*D trial, completed in 2006, was conducted to evaluate the effectiveness of current treatment approaches in patients with MDD. ^bRemission defined as QIDS-SR16 ≤5 (1-5=No depression, 6-10=Mild depression, 11-15=Moderate depression, 16-20=Severe depression, 21-27=Very severe depression) at exit. Patients were evaluated across 4 successive steps of therapy (each step consisted of a 12-week, open-label trial, with an additional 2 weeks for patients deemed close to remission), either switching or augmenting treatment if an adequate response was not achieved. Those who responded adequately at any step could enter a 12-month naturalistic follow-up phase. ^cAnalysis of claims of privately insured individuals from OptumHealth Care Solutions, July 2009-March 2015; retrospective, longitudinal, matched cohort design. ^dData were drawn from the 2013 US National Health and Wellness Survey (NHWS; N = 75,000). ^eSPRAVATO® was dosed twice weekly (56 mg on day 1; may be increased to 84 mg from day 4) from weeks 1-4, weekly (56 or 84 mg) from weeks 5-8, and weekly or Q2W (56 mg or 84 mg) from weeks 9-32. SPRAVATO® was given along with an OAD that elicited nonresponse at baseline. ^fQuetiapine XR was dosed daily, starting at 50 mg and titrated to 150 mg/day on day 3-4 or up to 300 mg/day on day 5 or after. Up-titration to >150 mg/day may be adapted based on individual tolerability and begin no later than the end of week 2. Quetiapine XR was then flexibly dosed (150-300 mg/day) from weeks 3-32. QUE XR was given along with an OAD that elicited nonresponse at baseline. ^gDue to SPRAVATO® needing to be administered under the supervision of a HCP, participants in the SPRAVATO® arm had twice weekly visits for the first 4 weeks of the study, while participants in the QUE XR arm were seen once weekly. A difference in the frequency of study visits between groups is a potential confounder. ^hRemission was defined as a MADRS total score of ≤10. ⁱThe full analysis set includes all randomly assigned patients. Percentages are based on the number of patients at each timepoint, using LOCF for missing data. Data for weeks 2 and 4 correspond to day 15 and 29, respectively.

AP, antipsychotic; CI, confidence interval; HCP, healthcare professional; HRU, healthcare resource utilization; LOCF, last observation carried forward; LS, least squares; MADRS, Montgomery-Asberg Depression Rating Scale; MDD, major depressive disorder; MDSI, major depressive disorder with suicidal ideation; NMDA, N-methyl D-aspartate; OAD, oral antidepressant; OR, odds ratio; Q2W, every 2 weeks; QUE XR, quetiapine extended release; REMS, risk evaluation and mitigation strategies; SE, standard error; SNRI, serotonin-norepinephrine reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor; TEAEs, treatment-emergent adverse events; TRD, treatment-resistant depression; USPI, US Prescribing Information; WPL, work productivity loss.

1. Rush AJ, et al. *Am J Psychiatry*. 2006;163(11):1905-1917. 2. Amos TB, et al. *J Clin Psychiatry*. 2018;79(2):17m11725. 3. Amos TB, et al. Poster Presented at: 29th Annual US Psychiatric & Mental Health Congress, October 21-24, 2016; San Antonio, Texas. 4. SPRAVATO® [Prescribing Information]. Titusville, NJ: Janssen Pharmaceuticals, Inc. 5. Reif A, et al. *N Engl J Med*. 2023;5:389(14):1298-1309. 6. McIntyre RS, et al. *CNS Spectr*. 2025;30(1):e26. 7. Mattingly G, et al. Poster presented at: Psych Congress Elevate; June 1-4, 2023; Las Vegas, Nevada. 8. Mattingly G, et al. Poster presented at NEI Congress. November 9-12, 2023. Colorado Springs, CO. 9. Clemens K, et al. *J Comp Eff Res*. 2025;14(7):e240092. 10. Clemens K, et al. *J Clin Psychiatry*. 2025;86(1):24m15425.