TREMFYA® (guselkumab) What is the dosing schedule of TREMFYA for active psoriatic arthritis?


Dosage form and strengths¹

Subcutaneous injection:



- 100 mg/mL in a single-dose One-Press patient-controlled injector
- 100 mg/mL in a single-dose prefilled syringe
- 100 mg/mL in a single-dose prefilled pen (TREMFYA PEN)

Recommended dosage schedule for adults with active psoriatic arthritis¹



Tremfya is administered as a single 100 mg injection under the skin at weeks 0 and 4, and then every 8 weeks.




If your healthcare provider decides that you or a caregiver may be able to give your injections of TREMFYA at home, **you should receive training** on the right way to prepare and inject TREMFYA before attempting to inject.



Do not try to inject yourself until you have been shown the right way to give the injections by your healthcare provider.

Recommended dosage schedule for children 6 years of age and older who also weigh at least 88 pounds (40 kg) with active psoriatic arthritis¹



Tremfya is administered as a single 100 mg injection under the skin at weeks 0 and 4, and then every 8 weeks.



TREMFYA should be administered by a **healthcare provider** or an **adult caregiver** who has been shown the right way to give the injections by your healthcare provider.

This information is provided ONLY in response to your request and is not intended as medical advice, to promote the use of our product, or suggest using it in any manner other than as described in the PRESCRIBING INFORMATION. All decisions regarding your medical treatment should be made with your healthcare professional(s). Please contact your healthcare professional(s) to discuss the information in this response.